

Percentage of S_1 families of different varieties segregating for certain mutants.

Varieties	Mutants											
	virescent seedling	albino seedling	yellow seedling	pale green seedling	glossy seedling	pale yellow seedling	striped seedling	defective endosperm	vivipary	liguleless seedling	lazy	dwarf
Yellow Boosman	2	5	5	5	11	4	3	18	4	1	0	0
Potch. Pearl	0	1	0	0	1	0	1	6	1	0	0	1
Potch. Pearl Early	0	6	0	0	1	0	3	2	0	0	0	0
Wisconsin (Dykema)	1	8	0	2	2	2	3	16	11	2	0	1
Potch. Pearl 14 row	0	3	0	1	1	0	2	3	3	1	0	3
Early King	1	1	0	3	6	2	3	13	2	1	0	0
White Boosman	2	8	0	9	2	3	4	9	0	0	0	1
Pale Boosman	3	7	1	8	16	4	4	10	3	1	0	1
Early Pearl (Mosterd)	0	1	1	7	0	4	4	9	1	0	0	1
Potch. Pearl Synthetic	0	0	0	0	0	1	0	4	1	0	0	0
Wisconsin (Evans)	1	7	1	11	0	7	3	12	4	2	0	1
Anveld Synthetic	0	0	0	0	3	0	2	4	0	0	0	0
Hickory King	0	3	0	1	6	1	3	13	0	1	0	0
Golden Beauty	0	4	5	5	15	9	3	14	2	0	n	0
Anveld	2	1	0	2	3	1	0	3	1	0	0	0
Uys	0	7	0	5	0	3	4	22	0	0	0	0
Peruvian	1	2	0	0	3	5	2	5	0	0	0	0
White Stalk Silver King	0	5	2	3	4	0	2	10	1	0	0	0
American White Flint	3	8	3	8	0	4	3	9	0	0	1	2
Natal 8 row	1	4	1	3	2	4	1	12	0	0	0	0
Mic Success	0	4	0	0	0	6	2	14	0	0	0	0
Hotnot	0	6	1	4	4	4	0	3	0	0	0	0
Teko Yellow	0	2	1	3	3	2	4	8	0	0	0	1
King's Cross	1	1	3	4	1	2	2	11	0	0	1	1
Salisbury White	1	2	0	4	3	4	3	12	0	0	0	0

G. W. Terblanche