

Table 2  
Distribution of Chromosome Numbers in the Microspores of 62-588-89

	Number of Chromosomes			
	10	11	12	13
Microspores observed	3	9	12	3
Microspores expected with random distribution	3	10	10	3

Raju S. K. Chaganti  
Uma S. Tantravahi

10. Intra- and intergenomic affinities of maize and Tripsacum chromosomes.

In order to assess the significance of chromosome association in maize-Tripsacum hybrids, intragenomic synaptic relationships of maize and Tripsacum were studied. The frequencies of chiasmatic associations and side by side associations of chromosomes (which probably represent homologous or homeologous pairing, Person, Canadian Jour. Bot. 33:11-30, 1955; Kimber and Riley, Bot. Rev. 29:480-531, 1963) at meta-anaphase of meiosis in a haploid maize plant and at metaphase in the haploid genome of Tripsacum from the triploid hybrid [(maize x *T. floridanum*) x maize] were scored. If the association frequency found in the hybrid (chiasmatic and side by side) resulted to a large extent from intergenomic pairing, then such association should be in excess of the sum of the individual pairing frequencies of the haploid genomes of the constituent species. If, on the other hand, the pairing in the hybrid is autosyndetic or predominantly so, then the association in the hybrid should be equal to or less than the sum of the pairing in the haploid genomes of the parental species. The mean per cell of chiasmatic association in haploid maize, haploid genome of Tripsacum, and the  $F_1$  hybrid was 0.06, 0.20, and 2.28 respectively while the mean per cell of side by side association in the three materials respectively was 0.28, 0.20, and 0.69. It can at once be seen that the pairing in the hybrid is much higher compared to the sum of mean pairing in the haploid genomes of maize and Tripsacum (2.28:0.26 chiasmatic and 0.69:0.48 side by side associations). Thus a significant amount of pairing in the maize-Tripsacum hybrids is intergenomic and involves maize and Tripsacum chromosomes.

Raju S. K. Chaganti

11. Nonrandom segregation of Tripsacum floridanum chromosomes in the progeny of the triploid hybrid [(maize x *T. floridanum*) x maize].

In order to study the mode of segregation of the Tripsacum chromosomes on the female side of the triploid hybrid, the distribution of chromosome numbers in a progeny population of 150 plants obtained by backcrossing the triploid hybrid by the maize parent was studied. The data are presented in the following table.

Random segregation of Tripsacum chromosomes would follow a distribution obtained by expanding the binomial  $(1/2 + 1/2)^{18}$ . The above data show clearly that the distribution is extremely skewed toward the side of the