

VII. AUTHOR INDEX TO VOLUME 48

| | |
|--|-----|
| Barratt, D. H. P., P. A. Peterson, and R. B. Flavell | 68 |
| Barratt, D. H. P., P. A. Peterson, and R. B. Flavell | 70 |
| Beckett, J. B. and M. G. Neuffer | 117 |
| Binder, W. D. and E. D. Styles. Assay for the effects of chemical and natural substances on growth using empty endo- spermic sac injection | 157 |
| Birchler, J. Preliminary genetic evidence that the <u>Adh₁</u> ^{FCm} duplication produces a single transcript | 54 |
| Bokde, S. Mutant "branched silkless" found in flint inbred . . . | 31 |
| Brewbaker, J. L. and S. H. Chang. Aphid resistance under apparent monogenic control | 37 |
| Brewbaker, J. L. and Y. Hasegawa. Nine maize peroxidase loci and their tissue specificities | 35 |
| Briggs, R. W. Cytoplasmic male sterility research: M ₂ generation from streptomycin treatments | 32 |
| Bryce, W. Amylo-1,6-glucosidase activity in maize endosperm . . . | 176 |
| Camussi, A., M. Motto and E. Ottaviano | 99 |
| Carlson, W. A mutant of chromosomal behavior in mitosis and meiosis | 76 |
| Carlson, W. B chromosomes induce nondisjunction in 9 B ⁹ pollen . | 81 |
| Ceska, O. and E. D. Styles. Expression of <u>pr</u> in plant tissues . . | 157 |
| Chandramouli. Hollow endosperm in maize | 11 |
| Chandramouli. Jointed seed syndrome in maize | 14 |
| Chandramouli. Induced dominant <u>Ce</u> (curled-entangled) | 15 |
| Chandravadana, P. and W. C. Galinat. Chromosome morphology of white multiple tester and its hybrid with Burnham Spreader . . | 94 |
| Chandravadana, P. and W. C. Galinat. More on affinity of <u>Tripsacum</u> chromosome 7 to maize and teosinte chromosome 4 . . . | 96 |
| Chandravadana, P., W. C. Galinat, and R. V. Tantravahi | 97 |
| Chang, S. H. and J. L. Brewbaker | 37 |
| Coe, E. H. Jr. Interactions of the brown-plant anthocyanin factors | 123 |
| Constantin, M. J. and L. M. Josephson | 148 |
| Crane, P. L. and T. R. Stierwalt | 139 |
| Cummins, J. E., D. B. Walden, and R. S. Verma | 163 |
| Dempsey, E. and M. M. Rhoades | 56 |

| | |
|---|-----|
| Doyle, G. G. Nonrandom pairing in trisomics with two standard chromosomes 3 and one chromosome 3 treated with ethyl methane sulfonate or nitrosoguanidine | 126 |
| Doyle, G. G. Telocentric 6L trisomes and their possible use in the commercial production of hybrid corn | 128 |
| Dunn, G. M., D. G. Routley, and B. J. Long | 131 |
| Filion, W. G. Differential Giemsa staining in maize | 150 |
| Flavell, R. B., D. H. P. Barratt, and P. A. Peterson | 68 |
| Flavell, R. B., D. H. P. Barratt, and P. A. Peterson | 70 |
| Foard, D., Y. Ma, and O. Nelson. The mutations, <u>de*-91</u> and <u>de*-92</u> | 169 |
| Forster, M. J. Heat induced autotetraploids of maize | 23 |
| Gabay, S. J. Growth of pollen from various inbred line and F ₁ sources on Cook and Walden basal medium | 43 |
| Gabay, S. J. and J. R. Laughnan | 38 |
| Gabay, S. J. and J. R. Laughnan. Sensitivity of pollen with Texas group male-sterile cytoplasm to <u>Helminthosporium maydis</u> race T pathotoxin | 42 |
| Gabay, S. J. and J. R. Laughnan. Linkage analysis in the male gametophyte | 44 |
| Galinat, W. C. A congruous background for the <u>tr</u> and <u>pd</u> genes | 93 |
| Galinat, W. C. On the individual breeder's gene pool | 93 |
| Galinat, W. C. On the relation of Tr9, a <u>Tripsacum</u> partial homeolog to M2, to maize MLO | 98 |
| Galinat, W. C. and P. Chandravadana | 94 |
| Galinat, W. C. and P. Chandravadana | 96 |
| Galinat, W. C., R. V. Tantravahi, and P. Chandravadana | 97 |
| Galli, G. and G. Gavazzi | 106 |
| Gavazzi, G. and G. Galli. Constitution of nonparental strands isolated from <u>R</u> intralocus recombinants | 106 |
| Gavazzi, G. and C. Piccardo. A new mutant affecting amino acid metabolism | 112 |
| Ghidoni, A. Nucleolar patterns in microspore quartets of trisomic 6, in relation to trisomic inheritance | 102 |
| Ghidoni, A. A dominant "blotch leaf" factor, homozygous lethal, located in chromosome 9 | 105 |
| Giles, K. L. Maize genetics studies by protoplast fusion | 29 |
| Gonella, J. and P. A. Peterson. Variegation associated with controlling element systems in tribal maize from Colombia | 66 |

| | |
|--|-----|
| Gracen, V. E. and C. O. Grogan. Evaluations of sources of cytoplasmic male sterility for use in multiplasm hybrid production | 20 |
| Greenblatt, I. M. Modulator as viewed through the pericarp | 186 |
| Greenblatt, I. M. Proximal-distal polarity of Modulator transpositions upon leaving the <u>P</u> locus | 188 |
| Greenblatt, I. M. Modulator: A modifier of crossing over | 189 |
| Greenblatt, I. M. Discussion of the above reports | 191 |
| Grogan, C. O. and V. E. Gracen | 20 |
| Gupta, D. and I. Kovacs. Cold tolerance of opaque-2 versus normal maize with and without fungicidal seed treatment | 4 |
| Hadjinov, M. I. and M. V. Tshumak. A simplified procedure of backcrossing for transferring the recessive trait to the recurrent parent | 89 |
| Hadjinov, M. I., K. I. Zima, A. A. Normov, and V. Z. Pakudyn. Combining ability of high protein opaque-2 maize lines for protein content in diallele crosses | 83 |
| Hasegawa, Y. and J. L. Brewbaker | 35 |
| Herczegh, M. Evaluation of heterozygous opaque hybrids of the Mv Syn A-3 O ₂ sublimes | 9 |
| Horn, J. D. Precise chromosome movements prior to somatic metaphase in maize | 165 |
| Horn, J. D. Cycloheximide-affected metaphase morphology in maize. | 167 |
| Horn, J. D. and D. B. Walden. The absence of nonhomologous associations of somatic chromosomes in maize | 165 |
| Iltis, Hugh H. Freezing the genetic landscape - the preservation of diversity in cultivated plants as an urgent social responsibility of the plant geneticist and plant taxonomist | 199 |
| Josephson, L. M. and M. J. Constantin. Attempts to modify cell organelles of Tcms by irradiation | 148 |
| Josephson, L. M. and H. C. Kincer. Tcms restoring genes in open-pollinated varieties | 146 |
| Josephson, L. M. and H. C. Kincer. Conversion of inbreds T115 and T202 to non-restoration of Tcms | 147 |
| Josephson, L. M. and H. C. Kincer. Conversion of virus-resistant inbreds to various male-sterile cytoplasms | 147 |
| Kermicle, J. Identical twins of dizygotic origin | 180 |
| Kincer, H. C. and L. M. Josephson | 146 |
| Kincer, H. C. and L. M. Josephson | 147 |
| Kovacs, I. A study of some characteristics of the heterozygous opaque-2 sister line crosses | 5 |

| | |
|--|-----|
| Kovacs, I. Comparison of some properties of the heterozygous opaque-2 hybrids and their normal analogues | 7 |
| Kovacs, I. and D. Gupta | 4 |
| Kyle, D. and E. D. Styles. The action of <u>Pl</u> in maize seedlings | 153 |
| Laughnan, J. R. and S. J. Gabay. Mutations that restore fertility to S male-sterile maize | 38 |
| Laughnan, J. R. and S. J. Gabay | 42 |
| Laughnan, J. R. and S. J. Gabay | 44 |
| Levings, C. S., III, and D. M. Shah. Comparison of mitochondrial DNA from hybrids with normal and Texas cytoplasms | 135 |
| Levites, E. V., S. G. Veprev, and S. I. Maletzky. Analysis of internode peroxidase in genetically determined dwarf forms of maize | 59 |
| Levites, E. V. and S. I. Maletzky. Study on ADH activity in the scutellum of diploid and tetraploid maize | 63 |
| Lin, B. Y. TB-10 breakpoints and marker genes on the long arm of chromosome 10 | 182 |
| Lin, B. Y. A seed-size effect associated with certain B-10 translocations | 184 |
| Liu, E. H. and J. G. Scandalios. Diaphorase isozyme patterns in the immature endosperm of <u>Zea mays</u> kernels | 142 |
| Loescher, W. H. and J. J. Mock. A nutrient medium for <u>in vitro</u> culture of maize and sorghum embryos | 64 |
| Long, B. J., G. M. Dunn, and D. G. Routley. Relation of hydroxamic acid content (DIMBOA) to resistance to <u>Helminthosporium turcicum</u> | 131 |
| Lorenzoni, C., M. Pozzi, and F. Salamini. Collapsed endosperm-1 (<u>cp₁</u>) location | 19 |
| Ma, Y. and O. Nelson. The endosperm proteins of <u>opaque-6</u> | 172 |
| Ma, Y., O. Nelson and D. Foard | 169 |
| Maguire, M. Apparent juxtaposition of homologues at premeiotic mitotic early anaphase | 149 |
| Maguire, M. Experimentally produced meiotic abnormalities | 149 |
| Maletzky, S. I. and E. V. Levites | 63 |
| Maletzky, S. I. and E. V. Poliakova | 58 |
| Maletzky, S. I., E. V. Levites and S. G. Veprev | 59 |
| Mock, J. J. and W. H. Loescher | 64 |
| Morgan, D. T. Jr. The <u>ws₃ lg₁ gl₂</u> region | 91 |
| Mottinger, J. P. Transposition of the control element affecting <u>bz-x3</u> | 193 |

| | |
|--|-----|
| Motto, M., E. Ottaviano, and A. Camussi | 99 |
| Nelson, O. and Y. Ma | 172 |
| Nelson, O., D. Foard, and Y. Ma | 169 |
| Neuffer, M. G. Absence of auxotrophic mutants in corn and other eukaryotes | 118 |
| Neuffer, M. G. Dominant mutants induced by EMS | 121 |
| Neuffer, M. G. Dominant disease lesion mutants | 121 |
| Neuffer, M. G. and J. B. Beckett. New mutants located with A-B translocations | 117 |
| Nielsen, G. and J. G. Scandalios. New alleles and chromosome localization of the locus for maize endopeptidase | 144 |
| Normov, A. A., V. Z. Pakudyn, M. I. Hadjinov, and K. I. Zima . . . | 83 |
| Ott, L. and J. G. Scandalios. Purification of maize peptidases . | 141 |
| Ottaviano, E., A. Camussi, and M. Motto. Selection of lines in relation to the effect of the opaque-2 gene on kernel weight in maize | 99 |
| Pakudyn, V. Z., M. I. Hadjinov, K. I. Zima, and A. A. Normov . . . | 83 |
| Peterson, P. A. Location of pg^m of the <u>En</u> system | 69 |
| Peterson, P. A. and J. Gonella | 66 |
| Peterson, P. A., R. B. Flavell, and D. H. P. Barratt. T-cytoplasm mitochondrial membrane activities | 68 |
| Peterson, P. A., R. B. Flavell, and D. H. P. Barratt. A sensitive assay for T mitochondria | 70 |
| Piccardo, C. and G. Gavazzi | 112 |
| Plewa, M. J. and D. F. Weber. Comparison of fatty acid percent- ages of diploid and monosomic <u>Zea mays</u> embryos | 46 |
| Poliakova, E. V. and S. I. Maletzky. Polymorphism of isozyme pattern of catalase in the endosperm of diploid maize strains . | 58 |
| Pozzi, M., F. Salamini, and C. Lorenzoni | 19 |
| Prior, C. L. and D. L. Shaver | 24 |
| Reddy, A. R. and G. M. Reddy. Confirmational studies of the accumulated flavonol in a_1 mutant aleurone tissue | 137 |
| Reddy, A. R. and G. M. Reddy. Differential action of intensifier gene | 138 |
| Reddy, A. R. and G. M. Reddy. Analysis of developing kernels . . | 138 |
| Reddy, A. R. and G. M. Reddy. Chemical analysis of certain mutant aleurone tissue | 139 |
| Reddy, G. M. and A. R. Reddy | 137 |
| Reddy, G. M. and A. R. Reddy | 138 |
| Reddy, G. M. and A. R. Reddy | 139 |

| | |
|--|-----|
| Rhoades, M. M. and E. Dempsey. Further studies on the cause of variation in frequency of chromatin loss induced by B chromosomes | 56 |
| Robertson, D. S. Linkage data for <u>yellow dwarf 2</u> on chromosome 3. | 70 |
| Robertson, D. S. Placement of <u>luteus*-1106</u> on the chromosome 4 linkage map | 72 |
| Robertson, D. S. Location of <u>pale yellow*-P11775 93</u> on chromosome 4 | 73 |
| Robertson, D. S. Linkage data for five pigment deficient mutants. | 75 |
| Routley, D. G., B. J. Long, and G. M. Dunn | 131 |
| Ryan, R. Preliminary studies of normal and male sterile cytoplasm in maize | 17 |
| Ryan, R. Effect of streptomycin on chlorophyll content in maize seedlings | 18 |
| Salamini, F., C. Lorenzoni, and M. Pozzi | 19 |
| Sartori, L. G bands in haploid maize | 19 |
| Scandalios, J. G. and E. H. Liu | 142 |
| Scandalios, J. G. and G. Nielson | 144 |
| Scandalios, J. G. and L. Ott | 141 |
| Scandalios, J. G. and J. C. Sorenson | 140 |
| Schwartz, D. On the quaternary structure of the temperature-sensitive mutant <u>Adh₁S-1100</u> | 52 |
| Shah, D. M. and C. S. Levings III | 135 |
| Shaver, D. L. Defective cytoplasm in <u>Zea</u> | 26 |
| Shaver, D. L. Inheritance of resistance to Sugar Cane Mosaic Virus (SCMV) in California | 27 |
| Shaver, D. L. and C. L. Prior. Cryptic earliness in maize | 24 |
| Shortess, D. K. Water-insoluble protein electrophoresis: a potential tool for maize gene product analysis | 132 |
| Singleton, W. R. Induction of mutants in maize pollen | 160 |
| Sorenson, J. C. and J. G. Scandalios. Partial purification of the catalase-specific inhibitor in maize | 140 |
| Stierwalt, T. R. and P. L. Crane. Soft endosperm genes | 139 |
| Styles, E. D. and W. D. Binder | 157 |
| Styles, E. D. and O. Ceska | 157 |
| Styles, E. D. and D. Kyle | 153 |
| Tantravahi, R. V., P. Chandravadana, and W. C. Galinat. Revision in a tentative identification of an extracted <u>Tripsacum</u> chromosome | 97 |

| | |
|---|-----|
| Ting, Y. C. Additional studies of the microsporocytes of Chalco (Mexico) teosintes | 16 |
| Ting, Y. C. Fine structure of the nucleolus of a diploid maize . | 17 |
| Tshumak, M. V. and M. I. Hadjinov | 89 |
| Veprev, S. G., S. I. Maletzky, and E. V. Levites | 59 |
| Verma, R. S., J. E. Cummins, and D. B. Walden. Chromosome aberrations produced by 5-bromodeoxyuridine (BUdR) and 5- fluorodeoxyuridine (FUdR) with concurrent exposure to U.V. in <u>Zea mays</u> L. root tips | 163 |
| Walden, D. B. and J. D. Horn | 165 |
| Walden, D. B., R. S. Verma, and J. E. Cummins | 163 |
| Ward, E. J. K10 and crossing over in Tp9Tp9 | 168 |
| Weber, D. F. A monosomic mapping method | 49 |
| Weber, D. F. and M. J. Flewa | 46 |
| Zima, K. I., A. A. Normov, V. Z. Pakudyn, and M. I. Hadjinov . . . | 83 |