

## VIII. RECENT MAIZE PUBLICATIONS

- Abedon, BG; Hatfield, RD; Tracy, WF, 2006. Cell wall composition in juvenile and adult leaves of maize (*Zea mays* L.). *J. Agric. Food Chem.* 54:3896-3900. Pubmed:16719512.
- Agrawal, GK; Rakwal, R, 2006. Rice proteomics: a cornerstone for cereal food crop proteomes. *Mass Spectrom. Rev.* 25:1-53. Pubmed:15957154.
- Ahmad, A; Wilde, GE; Whitworth, RJ; Zolnerowich, G, 2006. Effect of corn hybrids expressing the coleopteran-specific cry3Bb1 protein for corn rootworm control on above-ground insect predators. *J. Econ. Entomol.* 99:1085-1095. Pubmed:16937659.
- Ahmabadabi, M; Ruf, S; Bock, R, 2006. A leaf-based regeneration and transformation system for maize (*Zea mays* L.). *Transgenic Res.* (in press). Pubmed:17103238.
- Al-Abed, D; Rudrabhatla, S; Talla, R; Goldman, S, 2006. Split-seed: a new tool for maize researchers. *Planta* 223:1355-1360. Pubmed:16489455.
- Alleman, M; Sidorenko, L; McGinnis, K; Seshadri, V; Dorweiler, JE; White, J; Sikkink, K; Chandler, VL, 2006. An RNA-dependent RNA polymerase is required for paramutation in maize. *Nature* 442:295-298. Pubmed:16855589.
- Allerdings, E; Ralph, J; Steinhart, H; Bunzel, M, 2006. Isolation and structural identification of complex feruloylated heteroxylan side-chains from maize bran. *Phytochemistry* 67:1276-1286. Pubmed:16777153.
- Alvarez, S; Goodger, JQ; Marsh, EL; Chen, S; Asirvatham, VS; Schachtman, DP, 2006. Characterization of the maize xylem sap proteome. *J. Proteome Res.* 5:963-972. Pubmed:16602704.
- Alves, AP; Spencer, TA; Tabashnik, BE; Siegfried, BD, 2006. Inheritance of resistance to the Cry1Ab *Bacillus thuringiensis* toxin in *Ostrinia nubilalis* (Lepidoptera: Crambidae). *J. Econ. Entomol.* 99:494-501. Pubmed:16686152.
- Ambrus, H; Darko, E; Szabo, L; Bakos, F; Kiraly, Z; Barnabas, B, 2006. In vitro microspore selection in maize anther culture with oxidative-stress stimulators. *Protoplasma* 228:87-94. Pubmed:16937059.
- Amorim, EP; Oliveira-Amorim, VB; dos-Santos, JB; de-Souza, AP; de-Souza, JC, 2006. Genetic distance based on SSR and grain yield of inter and intrapopulational maize single cross hybrids. *Maydica* 51:507-513. Pubmed:20070123.
- Andersen, JR; Zein, I; Wenzel, G; Krutzfeldt, B; Eder, J; Ouzunova, M; Lubberstedt, T, 2007. High levels of linkage disequilibrium and associations with forage quality at a phenylalanine ammonia-lyase locus in European maize (*Zea mays* L.) inbreds. *Theor. Appl. Genet.* 114:307-319. Pubmed:17123062.
- Anderson, LK; Lai, A; Stack, SM; Rizzon, C; Gaut, BS, 2006. Uneven distribution of expressed sequence tag loci on maize pachytene chromosomes. *Genome Res.* 16:115-122. Pubmed:16339046.
- Andjelkovic, V; Thompson, R, 2006. Changes in gene expression in maize kernel in response to water and salt stress. *Plant Cell Rep.* 25:71-79. Pubmed:16362303.
- Andrieu, B; Hillier, J; Birch, C, 2006. Onset of sheath extension and duration of lamina extension are major determinants of the response of maize lamina length to plant density. *Ann. Bot. (Lond.)* 98:1005-1016. Pubmed:16926228.
- Anzala, F; Morere-Le Paven, MC; Fournier, S; Rondeau, D; Limami, AM, 2006. Physiological and molecular aspects of aspartate-derived amino acid metabolism during germination and post-germination growth in two maize genotypes differing in germination efficiency. *J. Exp. Bot.* 57:645-653. Pubmed:16415333.
- Asakura, Y; Barkan, A, 2006. *Arabidopsis* orthologs of maize chloroplast splicing factors promote splicing of orthologous and species-specific group II introns. *Plant Physiol.* 142:1656-1663. Pubmed:17071648.
- Babu, R; Nair, SK; Kumar, A; Rao, HS; Verma, P; Gahalain, A; Singh, IS; Gupta, HS, 2006. Mapping QTLs for popping ability in a popcorn x flint corn cross. *Theor. Appl. Genet.* 112:1392-1399. Pubmed:16525839.
- Badu-Apraku, B; Menkir, A, 2006. Registration of 16 extra-early maturing *Striga* resistant tropical maize inbred lines. *Crop Sci.* 46:1400-1401. Pubmed:20060918.
- Badu-Apraku, B; Menkir, A; Fakorede, MAB; Lum, AF; Obeng-Antwi, K, 2006. Multivariate analyses of the genetic diversity of forty-seven *Striga* resistant tropical early maturing maize inbred lines. *Maydica* 51:551-559. Pubmed:20070123.
- Badu-Apraku, B; Menkir, A; Kling, JG; Fakorede, MAB, 2006. Registration of 16 *Striga* resistant early maturing tropical maize inbred lines. *Crop Sci.* 46:1410-1411. Pubmed:20060918.
- Badu-Apraku, B; Twumasi-Afriyie, S; Sallah, PYK; Haag, W; Asiedu, E; Marfo, KA; Dapaah, S; Dzafi, BD, 2006. Registration of 'Obatanga GH' maize. *Crop Sci.* 46:1393-1395. Pubmed:20060918.
- Bai, L; Singh, M; Pitt, L; Sweeney, M; Brutnell, TP, 2007. Generating novel allelic variation through Activator insertional mutagenesis in maize. *Genetics* 175:981-992. Pubmed:17151236.
- Bai, QL; Chen, SJ; Dong, XL; Meng, QX; Yan, YL; Dai, JR, 2006. [Prediction of IVDMD with near infrared reflectance spectroscopy (NIRS) in maize stalk]. *Guang Pu Xue Yu FixGuang Pu Fen Xi* 26:271-274. Pubmed:16826904.
- Bai, WY; Zhao, LQ; Li, ZP; Xie, WQ; Zhao, YL; Li, CF, 2006. [Cloning and expression of cDNA for maize nonspecific lipid transfer protein as well as calmodulin-binding activity analysis of the expression product]. *Zhi Wu Sheng Li Yu Fen Zi Sheng Wu Xue Xue Bao* 32:570-576. Pubmed:17075181.
- Balint-Kurti, PJ; Blanco, M; Millard, M; Duvick, S; Holland, J; Clements, M; Holley, R; Carson, ML; Goodman, MM, 2006. Registration of 20 GEM maize breeding germplasm lines adapted to the southern USA. *Crop Sci.* 46:996-998. Pubmed:20060718.

- Barcaccia, G; Pallottini, L; Parrini, P; Lucchin, M, 2006. A genetic linkage map of a flint maize (*Zea mays* var. *indurata* L.) Italian landrace using a one-way pseudo-testcross strategy and multilocus PCR-based markers. *Maydica* 51:469-480. Pubmed:20070123.
- Barkan, A; Klipcan, L; Ostersetter, O; Kawamura, T; Asakura, Y; Watkins, KP, 2007. The CRM domain: an RNA binding module derived from an ancient ribosome-associated protein. *RNA* 13:55-64. Pubmed:17105995.
- Barker, CL; Baillie, BK; Hammond-Kosack, KE; Jones, JD; Jones, DA, 2006. Dominant-negative interference with defence signalling by truncation mutations of the tomato *Cf-9* disease resistance gene. *Plant J.* 46:385-399. Pubmed:16623900.
- Barker, CL; Talbot, SJ; Jones, JD; Jones, DA, 2006. A tomato mutant that shows stunting, wilting, progressive necrosis and constitutive expression of defence genes contains a recombinant *Hcr9* gene encoding an autoactive protein. *Plant J.* 46:369-384. Pubmed:16623899.
- Barbero, C; Muniz, LM; Gomez, E; Hueros, G; Royo, J, 2006. Molecular dissection of the interaction between the transcriptional activator *ZmMRP-1* and the promoter of *BETL-1*. *Plant Mol. Biol.* 62:655-668. Pubmed:17001496.
- Barret, P; Brinkman, M; Beckert, M, 2006. A sequence related to rice *Pong* transposable element displays transcriptional activation by in vitro culture and reveals somaclonal variations in maize. *Genome* 49:1399-1407. Pubmed:17426755.
- Barriere, Y; Alber, D; Dolstra, O; Lapierre, C; Motto, M; Ordas, A; Van-Waes, J; Vlasminikel, L; Welcker, C; Monod, JP, 2006. Past and prospects of forage maize breeding in Europe. II. History, germplasm evolution and correlative agronomic changes. *Maydica* 51:435-449. Pubmed:20070123.
- Barriere, Y; Denoue, D; Briand, M; Simon, M; Jouanin, L; Durand-Tardif, M, 2006. Genetic variations of cell wall digestibility related traits in floral stems of *Arabidopsis thaliana* accessions as a basis for the improvement of the feeding value in maize and forage plants. *Theor. Appl. Genet.* 113:163-175. Pubmed:16783597.
- Barry, JK; Selinger, DA; Wang, C; Olsen, OA; Rao, AG, 2006. Biochemical characterization of a truncated penta-EF-hand Ca<sup>2+</sup> binding protein from maize. *Biochim. Biophys. Acta* 1764:239-245. Pubmed:16269278.
- Bashir, K; Inoue, H; Nagasaka, S; Takahashi, M; Nakanishi, H; Mori, S; Nishizawa, NK, 2006. Cloning and characterization of deoxymugineic acid synthase genes from graminaceous plants. *J. Biol. Chem.* 281:32395-32402. Pubmed:16926158.
- Bastida, M; Graziano, E; Roca, R; Lopez, I; Sanchez-Pons, N; Puigdomenech, P; Vicent, CM, 2006. A maize defective-kernel mutant (*longcell*) characterized by tubular cells, severe morphological alterations and induction of cell death. *Planta* 223:755-768. Pubmed:16231157.
- Bauer, MJ; Birchler, JA, 2006. Organization of endoreduplicated chromosomes in the endosperm of *Zea mays* L. *Chromosoma* 115:383-394. Pubmed:16741707.
- Bertoia, L; Lopez, C; Burak, R, 2006. Biplot analysis of forage combining ability in maize landraces. *Crop Sci.* 46:1346-1353. Pubmed:20060918.
- Beyer, AJ; Wang, K; Umble, AN; Wolt, JD; Cunnick, JE, 2007. Low-dose exposure and immunogenicity of transgenic maize expressing the *Escherichia coli* heat-labile toxin B subunit. *Environ. Health Perspect.* 115:354-360. Pubmed:17431483.
- Bi, IV; McMullen, MD; Sanchez-Villeda, H; Schroeder, S; Gardiner, J; Polacco, M; Soderlund, C; Wing, R; Fang, Z; Coe, EH, Jr., 2006. Single nucleotide polymorphisms and insertion-deletions for genetic markers and anchoring the maize fingerprint contig physical map. *Crop Sci.* 46:12-21. Pubmed:20060524.
- Bianchi, A; Peterson, PA, 2006. Major Goodman: An Iowa farm youth devoted his career to broadening the maize genetic base. *Maydica* 51:1. Pubmed:20061017.
- Bird, RM, 2006. Zea models and matrices, measures and materials. *Maydica* 51:103-122. Pubmed:20061017.
- Blackburn, EH; Greider, CW; Szostak, JW, 2006. Telomeres and telomerase: the path from maize, *Tetrahymena* and yeast to human cancer and aging. *Nat. Med.* 12:1133-1138. Pubmed:17024208.
- Blanc, G; Charchosset, A; Mangin, B; Gallais, A; Moreau, L, 2006. Connected populations for detecting quantitative trait loci and testing for epistasis: an application in maize. *Theor. Appl. Genet.* 113:206-224. Pubmed:16791688.
- Bloom, AJ; Frensch, J; Taylor, AR, 2006. Influence of inorganic nitrogen and pH on the elongation of maize seminal roots. *Ann. Bot. (Lond.)* 97:867-873. Pubmed:16373369.
- Boddu, J; Jiang, C; Sangar, V; Olson, T; Peterson, T; Chopra, S, 2006. Comparative structural and functional characterization of sorghum and maize duplications containing orthologous myb transcription regulators of 3-deoxyflavonoid biosynthesis. *Plant Mol. Biol.* 60:185-199. Pubmed:16429259.
- Bolle, N; Kempken, F, 2006. Mono- and dicotyledonous plant-specific RNA editing sites are correctly edited in both in organello systems. *FEBS Lett.* 580:4443-4448. Pubmed:16857195.
- Bomblies, K; Doebley, JF, 2006. Pleiotropic effects of the duplicate maize *FLORICAULA/LEAFY* genes *zfl1* and *zfl2* on traits under selection during maize domestication. *Genetics* 172:519-531. Pubmed:16204211.
- Bondili, JS; Castilho, A; Mach, L; Glossl, J; Steinkellner, H; Altmann, F; Strasser, R, 2006. Molecular cloning and heterologous expression of beta1,2-xylosyltransferase and core alpha1,3-fucosyltransferase from maize. *Phytochemistry* 67:2215-2224. Pubmed:16920165.
- Bordes, J; Charmet, G; de Vaulx, RD; Pollacsek, M; Beckert, M; Gallais, A, 2006. Doubled haploid versus S1 family recurrent selection for testcross performance in a maize population. *Theor. Appl. Genet.* 112:1063-1072. Pubmed:16432736.
- Bortiri, E; Chuck, G; Vollbrecht, E; Rocheford, T; Martienssen, R; Hake, S, 2006. *ramosa2* encodes a LATERAL ORGAN BOUNDARY domain protein that determines the fate of stem cells in branch meristems of maize. *Plant Cell* 18:574-585. Pubmed:16399802.

- Bortiri, E; Jackson, D; Hake, S, 2006. Advances in maize genomics: the emergence of positional cloning. *Curr. Opin. Plant Biol.* 9:164-171. Pubmed:16458573.
- Bouchabke, O; Tardieu, F; Simonneau, T, 2006. Leaf growth and turgor in growing cells of maize (*Zea mays* L.) respond to evaporative demand under moderate irrigation but not in water-saturated soil. *Plant Cell Environ.* 29:1138-1148. Pubmed:17080939.
- Bouranis, DL; Chorianopoulou, SN; Kollias, C; Maniou, P; Protonotarios, VE; Siyannis, VF; Hawkesford, MJ, 2006. Dynamics of aerenchyma distribution in the cortex of sulfate-deprived adventitious roots of maize. *Ann. Bot. (Lond.)* 97:695-704. Pubmed:16481362.
- Boutte, Y; Crosnier, MT; Carraro, N; Traas, J; Satiat-Jeunemaitre, B, 2006. The plasma membrane recycling pathway and cell polarity in plants: studies on PIN proteins. *J. Cell Sci.* 119:1255-1265. Pubmed:16522683.
- Boyer, JS; McLaughlin, JE, 2007. Functional reversion to identify controlling genes in multigenic responses: analysis of floral abortion. *J. Exp. Bot.* 58:267-277. Pubmed:17105969.
- Brady, SM; Song, S; Dhugga, KS; Rafalski, JA; Benfey, PN, 2007. Combining expression and comparative evolutionary analysis. The COBRA gene family. *Plant Physiol.* 143:172-187. Pubmed:17098858.
- Braun, DM; Ma, Y; Inada, N; Muszynski, MG; Baker, RF, 2006. *tie-dyed1* regulates carbohydrate accumulation in maize leaves. *Plant Physiol.* 142:1511-1522. Pubmed:17071639.
- Brouquisse, R; Rolin, D; Cortes, S; Gaudillere, M; Evrard, A; Roby, C, 2007. A metabolic study of the regulation of proteolysis by sugars in maize root tips: effects of glycerol and dihydroxyacetone. *Planta* 225:693-709. Pubmed:16944197.
- Brown, PJ; Klein, PE; Bortiri, E; Acharya, CB; Rooney, WL; Kresovich, S, 2006. Inheritance of inflorescence architecture in sorghum. *Theor. Appl. Genet.* 113:931-942. Pubmed:16847662.
- Bruggmann, R; Bharti, AK; Gundlach, H; Lai, J; Young, S; Pontaroli, AC; Wei, F; Haberer, G; Fuks, G; Du, C; Raymond, C; Estep, MC; Liu, R; Bennetzen, JL; Chan, AP; Rabinowicz, PD; Quackenbush, J; Barbazuk, WB; Wing, RA; Birren, B; Nusbaum, C; Rounsley, S; Mayer, KF; Messing, J, 2006. Uneven chromosome contraction and expansion in the maize genome. *Genome Res.* 16:1241-1251. Pubmed:16902087.
- Brunk, CG, 2006. Public knowledge, public trust: understanding the 'knowledge deficit'. *Community Genet.* 9:178-183. Pubmed:16741347.
- Bubeck, DM; Carbone, MR; Fox, RL; Hoffbeck, MD; Segebart, RL; Stucker, DS, 2006. Breeding progress measured in eight elite inbred families. *Maydica* 51:141-149. Pubmed:20061017.
- Buckler, ES; Gaut, BS; McMullen, MD, 2006. Molecular and functional diversity of maize. *Curr. Opin. Plant Biol.* 9:172-176. Pubmed:16459128.
- Buckler, ES; Goodman, MM; Holtsford, TP; Doebley, JF; Sanchez-G, J, 2006. Phylogeography of the wild subspecies of *Zea mays*. *Maydica* 51:123-134. Pubmed:20061017.
- Burnell, JN; Chastain, CJ, 2006. Cloning and expression of maize-leaf pyruvate, Pi dikinase regulatory protein gene. *Biochem. Biophys. Res. Commun.* 345:675-680. Pubmed:16696949.
- Burza, AM; Pekala, I; Sikora, J; Siedlecki, P; Malagocki, P; Bucholc, M; Koper, L; Zielenkiewicz, P; Dadlez, M; Dobrowolska, G, 2006. *Nicotiana tabacum* osmotic stress-activated kinase is regulated by phosphorylation on Ser-154 and Ser-158 in the kinase activation loop. *J. Biol. Chem.* 281:34299-34311. Pubmed:16980311.
- Butron, A; Santiago, R; Mansilla, P; Pintos-Varela, C; Ordas, A; Malvar, RA, 2006. Maize (*Zea mays* L.) genetic factors for preventing fumonisin contamination. *J. Agric. Food Chem.* 54:6113-6117. Pubmed:16881725.
- Camoni, L; Marra, M; Garufi, A; Visconti, S; Aducci, P, 2006. The maize root plasma membrane H(+)-ATPase is regulated by a sugar-induced transduction pathway. *Plant Cell Physiol.* 47:743-747. Pubmed:16614095.
- Campos, H; Cooper, M; Edmeades, GO; Loffler, C; Schussler, JR; Ibanez, M, 2006. Changes in drought tolerance in maize associated with fifty years of breeding for yield in the US corn belt. *Maydica* 51:369-381. Pubmed:20061003.
- Camus-Kulandaivelu, L; Veyrieras, JB; Madur, D; Combes, V; Fourmann, M; Barraud, S; Dubreuil, P; Gouesnard, B; Manicacci, D; Charcosset, A, 2006. Maize adaptation to temperate climate: relationship between population structure and polymorphism in the *Dwarf8* gene. *Genetics* 172:2449-2463. Pubmed:16415370.
- Canaran, P; Stein, L; Ware, D, 2006. Look-Align: an interactive web-based multiple sequence alignment viewer with polymorphism analysis support. *Bioinformatics* 22:885-886. Pubmed:16473875.
- Cankar, K; Stebih, D; Dre, T; Zel, J; Gruden, K, 2006. Critical points of DNA quantification by real-time PCR--effects of DNA extraction method and sample matrix on quantification of genetically modified organisms. *BMC Biotechnol.* 6:37. Pubmed:16907967.
- Cardinal, AJ; Lee, M; Guthrie, WD; Bing, J; Austin, DF; Veldboom, LR; Senior, ML, 2006. Mapping of factors for resistance to leaf-blade feeding by European corn borer (*Ostrinia nubilalis*) in maize. *Maydica* 51:93-102. Pubmed:20061017.
- Carena, MJ; Wicks, ZW, III, 2006. Maize population hybrids: An exploitation of US temperate public genetic diversity in reserve. *Maydica* 51:201-208. Pubmed:20061003.
- Carlson, WR, 2006. Unstable inheritance of maize B-type chromosomes that lack centric heterochromatin. *Genome* 49:420-431. Pubmed:16767167.
- Carpane, P; Laguna, IG; Virla, E; Paradell, S; Murua, I; Gimenez-Pecchi, MP, 2006. Experimental transmission of corn stunt *Spiroplasma* present in different regions of Argentina. *Maydica* 51:461-468. Pubmed:20070123.

- Carraro, N; Forestan, C; Canova, S; Traas, J; Varotto, S, 2006. *ZmPIN1a* and *ZmPIN1b* encode two novel putative candidates for polar auxin transport and plant architecture determination of maize. *Plant Physiol.* 142:254-264. Pubmed:16844839.
- Carson, ML; Balint-Kurti, PJ; Blanco, M; Millard, M; Duvick, S; Holley, R; Hudyncia, J; Goodman, MM, 2006. Registration of nine high-yielding tropical by temperate maize germplasm lines adapted for the southern USA. *Crop Sci.* 46:1825-1826. Pubmed:20061017.
- Carson, ML; Goodman, MM, 2006. Pathogenicity, aggressiveness, and virulence of three species of *Cercospora* associated with gray leaf spot of maize. *Maydica* 51:89-92. Pubmed:20061017.
- Carvalho, AO; Souza-Filho, GA; Ferreira, BS; Branco, AT; Araujo, IS; Fernandes, KV; Retamal, CA; Gomes, VM, 2006. Cloning and characterization of a cowpea seed lipid transfer protein cDNA: expression analysis during seed development and under fungal and cold stresses in seedlings' tissues. *Plant Physiol. Biochem.* 44:732-742. Pubmed:17084637.
- Casati, P; Stapleton, AE; Blum, JE; Walbot, V, 2006. Genome-wide analysis of high-altitude maize and gene knockdown stocks implicates chromatin remodeling proteins in response to UV-B. *Plant J.* 46:613-627. Pubmed:16640598.
- Castells, E; Puigdomenech, P; Casacuberta, JM, 2006. Regulation of the kinase activity of the MIK GCK-like MAP4K by alternative splicing. *Plant Mol. Biol.* 61:747-756. Pubmed:16897489.
- Cavalar, M; Philippen, Y; Kreuzaler, F; Peterhansel, C, 2006. A drastic reduction in *DOF1* transcript levels does not affect C(4)-specific gene expression in maize. *J. Plant Physiol. (in press)*. Pubmed:17178169.
- Cerdeira, AL; Duke, SO, 2006. The current status and environmental impacts of glyphosate-resistant crops: a review. *J. Environ. Qual.* 35:1633-1658. Pubmed:16899736.
- Cervantes-Cervantes, M; Gallagher, CE; Zhu, C; Wurtzel, ET, 2006. Maize cDNAs expressed in endosperm encode functional farnesyl diphosphate synthase with geranylgeranyl diphosphate synthase activity. *Plant Physiol.* 141:220-231. Pubmed:16581875.
- Chan, AP; Pertea, G; Cheung, F; Lee, D; Zheng, L; Whitelaw, C; Pontaroli, AC; SanMiguel, P; Yuan, Y; Bennetzen, J; Barbazuk, WB; Quackenbush, J; Rabinowicz, PD, 2006. The TIGR Maize Database. *Nucl. Acids Res.* 34:D771-776. Pubmed:16381977.
- Che, P; Love, TM; Frame, BR; Wang, K; Carriquiry, AL; Howell, SH, 2006. Gene expression patterns during somatic embryo development and germination in maize Hi II callus cultures. *Plant Mol. Biol.* 62:1-14. Pubmed:16845483.
- Chilcutt, CF, 2006. Cannibalism of *Helicoverpa zea* (Lepidoptera: Noctuidae) from *Bacillus thuringiensis* (Bt) transgenic corn versus non-Bt corn. *J. Econ. Entomol.* 99:728-732. Pubmed:16813305.
- Chilcutt, CF; Odvody, GN; Correa, JC; Remmers, J; Parker, RD, 2006. Decreased whorl and ear damage in nine mon810 *Bacillus thuringiensis* (Bt)-transgenic corn hybrids compared with their non-Bt counterparts. *J. Econ. Entomol.* 99:2164-2170. Pubmed:17195689.
- Ching, A; Dhugga, KS; Appenzeller, L; Meeley, R; Bourett, TM; Howard, RJ; Rafalski, A, 2006. *Brittle stalk 2* encodes a putative glycosylphosphatidylinositol-anchored protein that affects mechanical strength of maize tissues by altering the composition and structure of secondary cell walls. *Planta* 224:1174-1184. Pubmed:16752131.
- Chourey, PS; Jain, M; Li, QB; Carlson, SJ, 2006. Genetic control of cell wall invertases in developing endosperm of maize. *Planta* 223:159-167. Pubmed:16025339.
- Christou, P; Capell, T; Kohli, A; Gatehouse, JA; Gatehouse, AM, 2006. Recent developments and future prospects in insect pest control in transgenic crops. *Trends Plant Sci.* 11:302-308. Pubmed:16690346.
- Chu, CG; Faris, JD; Friesen, TL; Xu, SS, 2006. Molecular mapping of hybrid necrosis genes *Ne1* and *Ne2* in hexaploid wheat using microsatellite markers. *Theor. Appl. Genet.* 112:1374-1381. Pubmed:16518615.
- Chumakov, MI; Rozhok, NA; Velikov, VA; Tyrnov, VS; Volokhina, IV, 2006. [In planta transformation of maize through inoculation of *Agrobacterium* into the silks]. *Genetika* 42:1083-1088. Pubmed:17025158.
- Cistue, L; Soriano, M; Castillo, AM; Valles, MP; Sanz, JM; Echavarri, B, 2006. Production of doubled haploids in durum wheat (*Triticum turgidum* L.) through isolated microspore culture. *Plant Cell Rep.* 25:257-264. Pubmed:16220343.
- Clark, PL; Vaughn, TT; Meinke, LJ; Molina-Ochoa, J; Foster, JE, 2006. *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae) larval feeding behavior on transgenic maize (MON 863) and its isolate. *J. Econ. Entomol.* 99:722-727. Pubmed:16813304.
- Clark, RM; Wagler, TN; Quijada, P; Doebley, J, 2006. A distant upstream enhancer at the maize domestication gene *tb1* has pleiotropic effects on plant and inflorescent architecture. *Nat. Genet.* 38:594-597. Pubmed:16642024.
- Cleveland, DA; Soleri, D; Cuevas, FA; Crossa, J; Gepts, P, 2005. Detecting (trans)gene flow to landraces in centers of crop origin: lessons from the case of maize in Mexico. *Environ. Biosafety Res.* 4:197-208; discussion 209-115. Pubmed:16827547.
- Clough, RC; Pappu, K; Thompson, K; Beifuss, K; Lane, J; Delaney, DE; Harkey, R; Drees, C; Howard, JA; Hood, EE, 2006. Manganese peroxidase from the white-rot fungus *Phanerochaete chrysosporium* is enzymatically active and accumulates to high levels in transgenic maize seed. *Plant Biotechnol. J.* 4:53-62. Pubmed:17177785.
- Coe, EH; Schaeffer, ML, 2006. Uncaging mutants: Moving from menageries to menages. *Maydica* 51:263-267. Pubmed:20061003.
- Colasanti, J; Tremblay, R; Wong, AY; Coneva, V; Kozaki, A; Mable, BK, 2006. The maize *INDETERMINATE1* flowering time regulator defines a highly conserved zinc finger protein family in higher plants. *BMC Genomics* 7:158. Pubmed:16784536.
- Cona, A; Rea, G; Botta, M; Corelli, F; Federico, R; Angelini, R, 2006. Flavin-containing polyamine oxidase is a hydrogen peroxide source in the oxidative response to the protein phosphatase inhibitor cantharidin in *Zea mays* L. *J. Exp. Bot.* 57:2277-2289. Pubmed:16831849.
- Coque, M; Gallais, A, 2006. Genomic regions involved in response to grain yield selection at high and low nitrogen fertilization in maize. *Theor. Appl. Genet.* 112:1205-1220. Pubmed:16552555.

- Creff, A; Leonard, B; Desnos, T, 2006. Targeted *Ds*-tagging strategy generates high allelic diversity at the *Arabidopsis HY2* locus. *Plant Mol. Biol.* 61:603-613. Pubmed:16897478.
- Crimi, M; Astegno, A; Zoccatelli, G; Esposito, MD, 2006. Pro-apoptotic effect of maize lipid transfer protein on mammalian mitochondria. *Arch. Biochem. Biophys.* 445:65-71. Pubmed:16325144.
- Crowder, DW; Onstad, DW; Gray, ME, 2006. Planting transgenic insecticidal corn based on economic thresholds: consequences for integrated pest management and insect resistance management. *J. Econ. Entomol.* 99:899-907. Pubmed:16813328.
- D'Andrea, KE; Otegui, ME; Cirillo, AG; Eyherabide, G, 2006. Genotypic variability in morphological and physiological traits among maize inbred lines-nitrogen responses. *Crop Sci.* 46:1266-1276. Pubmed:20060918.
- da Silva, AL; Sperling, P; Horst, W; Franke, S; Ott, C; Becker, D; Stass, A; Lorz, H; Heinz, E, 2006. A possible role of sphingolipids in the aluminium resistance of yeast and maize. *J. Plant Physiol.* 163:26-38. Pubmed:16360801.
- Darie, CC; De Pascalis, L; Muttschler, B; Haehnel, W, 2006. Studies of the Ndh complex and photosystem II from mesophyll and bundle sheath chloroplasts of the C4-type plant *Zea mays*. *J. Plant Physiol.* 163:800-808. Pubmed:16436305.
- Dean, JM; De Moraes, CM, 2006. Effects of genetic modification on herbivore-induced volatiles from maize. *J. Chem. Ecol.* 32:713-724. Pubmed:16718567.
- Detarsio, E; Alvarez, CE; Saigo, M; Andreo, CS; Drincovich, MF, 2007. Identification of domains involved in tetramerization and malate inhibition of maize C4-NADP-malic enzyme. *J. Biol. Chem.* 282:6053-6060. Pubmed:17150960.
- Devos, Y; Reheul, D; Thas, O; De Clercq, EM; Cordemans, K, 2006. Spatial impact of isolation distances between parcels of GM and non-GM maize. *Commun. Agric. Appl. Biol. Sci.* 71:25-28. Pubmed:17191467.
- Diao, X; Freeling, M; Lisch, D, 2006. Horizontal transfer of a plant transposon. *PLoS Biol.* 4:e5. Pubmed:16336045.
- Diao, XM; Lisch, D, 2006. *Mutator* transposon in maize and MULEs in the plant genome. *Yi Chuan Xue Bao* 33:477-487. Pubmed:16800377.
- Dicko, MH; Gruppen, H; Hilhorst, R; Voragen, AG; van Berkel, WJ, 2006. Biochemical characterization of the major sorghum grain peroxidase. *FEBS J.* 273:2293-2307. Pubmed:16650004.
- Dinelli, G; Bonetti, A; Marotti, I; Minelli, M; Navarrete-Casas, M; Segura-Carretero, A; Fernandez-Gutierrez, A, 2006. Quantitative-competitive polymerase chain reaction coupled with slab gel and capillary electrophoresis for the detection of roundup ready soybean and maize. *Electrophoresis* 27:4029-4038. Pubmed:16983630.
- Ding, SY; Himmel, ME, 2006. The maize primary cell wall microfibril: a new model derived from direct visualization. *J. Agric. Food Chem.* 54:597-606. Pubmed:16448156.
- Djakovic, S; Dyachok, J; Burke, M; Frank, MJ; Smith, LG, 2006. BRICK1/HSPC300 functions with SCAR and the ARP2/3 complex to regulate epidermal cell shape in *Arabidopsis*. *Development* 133:1091-1100. Pubmed:16481352.
- Djukanovic, V; Orczyk, W; Gao, H; Sun, X; Garrett, N; Zhen, S; Gordon-Kamm, W; Barton, J; Lyznik, LA, 2006. Gene conversion in transgenic maize plants expressing FLP/FRT and Cre/loxP site-specific recombination systems. *Plant Biotechnol. J.* 4:345-357. Pubmed:17147640.
- Dlaskova, A; Spacek, T; Skobisova, E; Santorova, J; Jezek, P, 2006. Certain aspects of uncoupling due to mitochondrial uncoupling proteins in vitro and in vivo. *Biochim. Biophys. Acta* 1757:467-473. Pubmed:16781660.
- Doebley, JF; Gaut, BS; Smith, BD, 2006. The molecular genetics of crop domestication. *Cell* 127:1309-1321. Pubmed:17190597.
- Dong, YB; Li, YL; Niu, SZ, 2006. [QTL analysis of starch content in maize kernels using the trisomic inheritance of the endosperm model]. *Yi Chuan* 28:1401-1406. Pubmed:17098709.
- Doust, AN; Kellogg, EA, 2006. Effect of genotype and environment on branching in weedy green millet (*Setaria viridis*) and domesticated foxtail millet (*Setaria italica*) (Poaceae). *Mol. Ecol.* 15:1335-1349. Pubmed:16626457.
- Dowd, PF; Holmes, RA; Pinkerton, TS; Johnson, ET; Lagrimini, LM; Boston, RS, 2006. Relative activity of a tobacco hybrid expressing high levels of a tobacco anionic peroxidase and maize ribosome-inactivating protein against *Helicoverpa zea* and *Lasiodesma serricornis*. *J. Agric. Food Chem.* 54:2629-2634. Pubmed:16569054.
- Dresselhaus, T; Srilunchang, KO; Leljak-Levanic, D; Schreiber, DN; Garg, P, 2006. The fertilization-induced DNA replication factor MCM6 of maize shuttles between cytoplasm and nucleus, and is essential for plant growth and development. *Plant Physiol.* 140:512-527. Pubmed:16407440.
- Driscoll, SP; Prins, A; Olmos, E; Kunert, KJ; Foyer, CH, 2006. Specification of adaxial and abaxial stomata, epidermal structure and photosynthesis to CO<sub>2</sub> enrichment in maize leaves. *J. Exp. Bot.* 57:381-390. Pubmed:16371401.
- Drozak, A; Romanowska, E, 2006. Acclimation of mesophyll and bundle sheath chloroplasts of maize to different irradiances during growth. *Biochim. Biophys. Acta* 1757:1539-1546. Pubmed:17034754.
- Du, C; Swigonova, Z; Messing, J, 2006. Retrotranspositions in orthologous regions of closely related grass species. *BMC Evol. Biol.* 6:62. Pubmed:16914031.
- Du, J; Zhu, Z; Li, WC, 2006. Over-expression of exotic superoxide dismutase gene *MnSOD* and increase in stress resistance in maize. *Zhi Wu Sheng Li Yu Fen Zi Sheng Wu Xue Bao* 32:57-63. Pubmed:16477132.
- Dubreuil, P; Warburton, M; Chastanet, M; Hoisington, D; Charcosset, A, 2006. More on the introduction of temperate maize into Europe: Large-scale bulk SSR genotyping and new historical elements. *Maydica* 51:281-291. Pubmed:20061003.
- Dumas, C; Gaude, T, 2006. Fertilization in plants: is calcium a key player? *Semin. Cell Dev. Biol.* 17:244-253. Pubmed:16563816.

- Duncan, KA; Hardin, SC; Huber, SC, 2006. The three maize sucrose synthase isoforms differ in distribution, localization, and phosphorylation. *Plant Cell Physiol.* 47:959-971. Pubmed:16760218.
- Duvick, SA; Pollak, LM; Edwards, JW; White, PJ, 2006. Altering the fatty acid composition of Corn Belt corn through *Tripsacum* introgression. *Maydica* 51:409-416. Pubmed:20061003.
- Echarte, L; Tollenaar, M, 2006. Kernel set in maize hybrids and their inbred lines exposed to stress. *Crop Sci.* 46:870-878. Pubmed:20060718.
- Edelmann, HG; Roth, U, 2006. Gravitropic plant growth regulation and ethylene: an unsought cardinal coordinate for a disused model. *Protoplasma* 229:183-191. Pubmed:17180500.
- Eichhorn, H; Lessing, F; Winterberg, B; Schirawski, J; Kamper, J; Muller, P; Kahmann, R, 2006. A ferroxidation/permeation iron uptake system is required for virulence in *Ustilago maydis*. *Plant Cell* 18:3332-3345. Pubmed:17138696.
- Eizaguirre, M; Albajes, R; Lopez, C; Eras, J; Lumbierres, B; Pons, X, 2006. Six years after the commercial introduction of Bt maize in Spain: field evaluation, impact and future prospects. *Transgenic Res.* 15:1-12. Pubmed:16475005.
- Emelyanov, A; Gao, Y; Naqvi, NI; Parinov, S, 2006. Trans-kingdom transposition of the maize dissociation element. *Genetics* 174:1095-1104. Pubmed:16951067.
- Emrich, SJ; Barbazuk, WB; Li, L; Schnable, PS, 2007. Gene discovery and annotation using LCM-454 transcriptome sequencing. *Genome Res.* 17:69-73. Pubmed:17095711.
- Emrich, SJ; Li, L; Wen, TJ; Yandea-Nelson, MD; Fu, Y; Guo, L; Chou, HH; Aluru, S; Ashlock, DA; Schnable, PS, 2007. Nearly identical paralogs: implications for maize (*Zea mays* L.) genome evolution. *Genetics* 175:429-439. Pubmed:17110490.
- Enoki, H; Miki, K; Koinuma, K, 2006. Mapping of quantitative trait loci associated with early flowering of a northern flint maize (*Zea mays* L.) inbred line. *Maydica* 51:515-523. Pubmed:20070123.
- Ermolli, M; Prospero, A; Balla, B; Querci, M; Mazzeo, A; Van Den Eede, G, 2006. Development of an innovative immunoassay for CP4EPSPS and Cry1AB genetically modified protein detection and quantification. *Food Addit. Contam.* 23:876-882. Pubmed:16901856.
- Eubanks, MW, 2006. A genetic bridge to utilize *Tripsacum* germplasm in maize improvement. *Maydica* 51:315-327. Pubmed:20061003.
- Falke, KC; Melchinger, AE; Flachenecker, C; Kusterer, B; Frisch, M, 2006. Comparison of linkage maps from F2 and three times intermated generations in two populations of European flint maize (*Zea mays* L.). *Theor. Appl. Genet.* 113:857-866. Pubmed:16832645.
- Fan, J; Wang, D; Liang, Z; Guo, M; Teng, M; Niu, L, 2006. Maize uroporphyrinogen III methyltransferase: overexpression of the functional gene fragments in *Escherichia coli* and one-step purification. *Protein Expr. Purif.* 46:40-46. Pubmed:16289918.
- Fan, L; Linker, R; Gepstein, S; Tanimoto, E; Yamamoto, R; Neumann, PM, 2006. Progressive inhibition by water deficit of cell wall extensibility and growth along the elongation zone of maize roots is related to increased lignin metabolism and progressive stelar accumulation of wall phenolics. *Plant Physiol.* 140:603-612. Pubmed:16384904.
- Fang, ZD; Laskey, JG; Huang, S; Bilyeu, KD; Morris, RO; Schmidt, FJ; English, JT, 2006. Combinatorially selected defense peptides protect plant roots from pathogen infection. *Proc. Natl. Acad. Sci. USA* 103:18444-18449. Pubmed:17030803.
- Farage, PK; Blowers, D; Long, SP; Baker, NR, 2006. Low growth temperatures modify the efficiency of light use by photosystem II for CO<sub>2</sub> assimilation in leaves of two chilling-tolerant C4 species, *Cyperus longus* L. and *Miscanthus x giganteus*. *Plant Cell Environ.* 29:720-728. Pubmed:17080621.
- Favarro, RD; Borecky, J; Colombi, D; Maia, IG, 2006. ZmPUMP encodes a fully functional monocot plant uncoupling mitochondrial protein whose affinity to fatty acid is increased with the introduction of a His pair at the second matrix loop. *Biochem. Biophys. Res. Commun.* 344:194-199. Pubmed:16603126.
- Fawcett, JA; Kawahara, T; Watanabe, H; Yasui, Y, 2006. A SINE family widely distributed in the plant kingdom and its evolutionary history. *Plant Mol. Biol.* 61:505-514. Pubmed:16830182.
- Feller, A; Hernandez, JM; Grotewold, E, 2006. An ACT-like domain participates in the dimerization of several plant basic-helix-loop-helix transcription factors. *J. Biol. Chem.* 281:28964-28974. Pubmed:16867983.
- Feltus, FA; Singh, HP; Lohithaswa, HC; Schulze, SR; Silva, TD; Paterson, AH, 2006. A comparative genomics strategy for targeted discovery of single-nucleotide polymorphisms and conserved-noncoding sequences in orphan crops. *Plant Physiol.* 140:1183-1191. Pubmed:16607031.
- Feng, L; Sebastian, S; Smith, S; Cooper, M, 2006. Temporal trends in SSR allele frequencies associated with long-term selection for yield of maize. *Maydica* 51:293-300. Pubmed:20061003.
- Ferreira, BS; Branco, AT; de Oliveira, MA; Pereira, MG; de Souza Filho, GA, 2006. Methodological improvements on extraction of nuclear proteins and its preliminary analysis during the maize (*Zea mays* L.) endosperm development. *Protein Pept. Lett.* 13:981-984. Pubmed:17168819.
- Ferreira Junior, JR; Ramos, AS; Chambergo, FS; Stambuk, BU; Muschellack, LK; Schumacher, R; El-Dorry, H, 2006. Functional expression of the maize mitochondrial URF13 down-regulates galactose-induced *GAL1* gene expression in *Saccharomyces cerevisiae*. *Biochem. Biophys. Res. Commun.* 339:30-36. Pubmed:16297867.
- Firbank, LG; Rothery, P; May, MJ; Clark, SJ; Scott, RJ; Stuart, RC; Boffey, CW; Brooks, DR; Champion, GT; Haughton, AJ; Hawes, C; Heard, MS; Dewar, AM; Perry, JN; Squire, GR, 2006. Effects of genetically modified herbicide-tolerant cropping systems on weed seedbanks in two years of following crops. *Biol. Lett.* 2:140-143. Pubmed:17148348.

- Flachenecker, C; Frisch, M; Falke, KC; Melchinger, AE, 2006. Genetic drift and selection effects of modified recurrent full-sib selection programs in two F2 populations of European flint maize. *Theor. Appl. Genet.* 113:1113-1120. Pubmed:16896708.
- Flachenecker, C; Frisch, M; Falke, KC; Melchinger, AE, 2006. Trends in population parameters and best linear unbiased prediction of progeny performance in a European F(2) maize population under modified recurrent full-sib selection. *Theor. Appl. Genet.* 112:483-491. Pubmed:16344984.
- Flor-Parra, I; Vranes, M; Kamper, J; Perez-Martin, J, 2006. Biz1, a zinc finger protein required for plant invasion by *Ustilago maydis*, regulates the levels of a mitotic cyclin. *Plant Cell* 18:2369-2387. Pubmed:16905655.
- Folcher, L; Eychenne, N; Weissenberger, A; Jarry, M; Regnault-Roger, C; Delos, M, 2006. Study of effects of Bt maize (*Zea mays*) events on Lepidoptera *Ostrinia nubilalis*, *Sesamia nonagrioides* in southwestern France. *Commun. Agric. Appl. Biol. Sci.* 71:227-231. Pubmed:17390797.
- Fornale, S; Sonbol, FM; Maes, T; Capellades, M; Puigdomenech, P; Rigau, J; Caparros-Ruiz, D, 2006. Down-regulation of the maize and *Arabidopsis thaliana* caffeic acid O-methyl-transferase genes by two new maize R2R3-MYB transcription factors. *Plant Mol. Biol.* 62:809-823. Pubmed:16941210.
- Frame, BR; McMurray, JM; Fonger, TM; Main, ML; Taylor, KW; Torney, FJ; Paz, MM; Wang, K, 2006. Improved *Agrobacterium*-mediated transformation of three maize inbred lines using MS salts. *Plant Cell Rep.* 25:1024-1034. Pubmed:16710703.
- Frame, BR; Paque, T; Wang, K, 2006. Maize (*Zea mays* L.). *Methods Mol. Biol.* 343:185-199. Pubmed:16988344.
- Franco, J; Crossa, J; Warburton, M-L; Taba, S, 2006. Sampling strategies for conserving maize diversity when forming core subsets using genetic markers. *Crop Sci.* 46:854-864. Pubmed:20060718.
- Frisch, M; Melchinger, AE, 2006. Marker-based prediction of the parental genome contribution to inbred lines derived from biparental crosses. *Genetics* 174:795-803. Pubmed:16888341.
- Fu, D; Huang, B; Xiao, Y; Muthukrishnan, S; Liang, GH, 2007. Overexpression of barley *hva1* gene in creeping bentgrass for improving drought tolerance. *Plant Cell Rep.* 26:467-477. Pubmed:17106681.
- Fu, S; Rogowsky, P; Nover, L; Scanlon, MJ, 2006. The maize heat shock factor-binding protein paralogs EMP2 and HSBP2 interact non-redundantly with specific heat shock factors. *Planta* 224:42-52. Pubmed:16331466.
- Fu, Y; Wen, TJ; Ronin, YI; Chen, HD; Guo, L; Mester, DI; Yang, Y; Lee, M; Korol, AB; Ashlock, DA; Schnable, PS, 2006. Genetic dissection of intermated recombinant inbred lines using a new genetic map of maize. *Genetics* 174:1671-1683. Pubmed:16951074.
- Fukayama, H; Tamai, T; Taniguchi, Y; Sullivan, S; Miyao, M; Nimmo, HG, 2006. Characterization and functional analysis of phosphoenolpyruvate carboxylase kinase genes in rice. *Plant J.* 47:258-268. Pubmed:16762031.
- Gallais, A; Moreau, L; Charcosset, A, 2007. Detection of marker-QTL associations by studying change in marker frequencies with selection. *Theor. Appl. Genet.* 114:669-681. Pubmed:17165081.
- Gambari, R; Feriotti, G, 2006. Surface plasmon resonance for detection of genetically modified organisms in the food supply. *J. AOAC Int.* 89:893-897. Pubmed:16792091.
- Gao, C; Hu, J; Zheng, Y; Zhang, S, 2006. [Antioxidant enzyme activities and proline content in maize seedling and their relationships to cold endurance]. *Ying Yong Sheng Tai Xue Bao* 17:1045-1050. Pubmed:16964938.
- Gardner, HD; Williams, WP; Windham, GL, 2006. Effects of Xenia on *Aspergillus flavus* infection and aflatoxin accumulation in maize inbreds. *Crop Sci.* 46:2151-2154. Pubmed:20061219.
- Gaspar, L; Sarvari, E; Morales, F; Szigeti, Z, 2006. Presence of 'PSI free' LHCI and monomeric LHCII and subsequent effects on fluorescence characteristics in lincomycin treated maize. *Planta* 223:1047-1057. Pubmed:16292567.
- Gathmann, A; Wirooks, L; Eckert, J; Schuphan, I, 2006. Spatial distribution of *Aglais urticae* (L.) and its host plant *Urtica dioica* (L.) in an agricultural landscape: implications for Bt maize risk assessment and post-market monitoring. *Environ. Biosafety Res.* 5:27-36. Pubmed:16978572.
- Gathmann, A; Wirooks, L; Hothorn, LA; Bartsch, D; Schuphan, I, 2006. Impact of Bt maize pollen (MON810) on lepidopteran larvae living on accompanying weeds. *Mol. Ecol.* 15:2677-2685. Pubmed:16842436.
- Gershater, M; Sharples, K; Edwards, R, 2006. Carboxylesterase activities toward pesticide esters in crops and weeds. *Phytochemistry* 67:2561-2567. Pubmed:17078983.
- Gibbons, DW; Bohan, DA; Rothery, P; Stuart, RC; Haughton, AJ; Scott, RJ; Wilson, JD; Perry, JN; Clark, SJ; Dawson, RJ; Firbank, LG, 2006. Weed seed resources for birds in fields with contrasting conventional and genetically modified herbicide-tolerant crops. *Proc. Biol. Sci.* 273:1921-1928. Pubmed:16822753.
- Gissot, L; Polge, C; Jossier, M; Girin, T; Bouly, JP; Kreis, M; Thomas, M, 2006. AKIN $\beta$  contributes to SnRK1 heterotrimeric complexes and interacts with two proteins implicated in plant pathogen resistance through its KIS/GBD sequence. *Plant Physiol.* 142:931-944. Pubmed:17028154.
- Glaring, MA; Koch, CB; Blennow, A, 2006. Genotype-specific spatial distribution of starch molecules in the starch granule: a combined CLSM and SEM approach. *Biomacromolecules* 7:2310-2320. Pubmed:16903676.
- Golubovskaya, IN; Hamant, O; Timofejeva, L; Wang, CJ; Braun, D; Meeley, R; Cande, WZ, 2006. Alleles of *afd1* dissect REC8 functions during meiotic prophase I. *J. Cell Sci.* 119:3306-3315. Pubmed:16868028.
- Gomez, E; Schulman, AH; Martinez-Izquierdo, JA; Vicent, CM, 2006. Integrase diversity and transcription of the maize retrotransposon Grande. *Genome* 49:558-562. Pubmed:16767181.

- Gonzalez, G; Comas, C; Confalonieri, V; Naranjo, CA; Poggio, L, 2006. Genomic affinities between maize and *Zea perennis* using classical and molecular cytogenetic methods (GISH-FISH). Chromosome Res. 14:629-635. Pubmed:16964569.
- Gonzalo, M; Vyn, TJ; Holland, JB; McIntyre, LM, 2006. Mapping density response in maize: a direct approach for testing genotype and treatment interactions. Genetics 173:331-348. Pubmed:16489238.
- Gorantla, M; Babu, PR; Lachagari, VB; Reddy, AM; Wusirika, R; Bennetzen, JL; Reddy, AR, 2007. Identification of stress-responsive genes in an indica rice (*Oryza sativa* L.) using ESTs generated from drought-stressed seedlings. J. Exp. Bot. 58:253-265. Pubmed:17132712.
- Gowda, M; Li, H; Alessi, J; Chen, F; Pratt, R; Wang, GL, 2006. Robust analysis of 5'-transcript ends (5'-RATE): a novel technique for transcriptome analysis and genome annotation. Nucl. Acids Res. 34:e126. Pubmed:17012272.
- Grant, WF; Owens, ET, 2006. *Zea mays* assays of chemical/radiation genotoxicity for the study of environmental mutagens. Mutat. Res. 613:17-64. Pubmed:16828334.
- Gray-Mitsumune, M; Matton, DP, 2006. The *Egg apparatus 1* gene from maize is a member of a large gene family found in both monocots and dicots. Planta 223:618-625. Pubmed:16341706.
- Greenway, H; Armstrong, W; Colmer, TD, 2006. Conditions leading to high CO<sub>2</sub> (>5 kPa) in waterlogged-flooded soils and possible effects on root growth and metabolism. Ann. Bot. (Lond.) 98:9-32. Pubmed:16644893.
- Griffiths, BS; Caul, S; Thompson, J; Birch, AN; Scrimgeour, C; Cortet, J; Foggo, A; Hackett, CA; Krogh, PH, 2006. Soil microbial and faunal community responses to bt maize and insecticide in two soils. J. Environ. Qual. 35:734-741. Pubmed:16585615.
- Gruber, MY; Wang, S; Ethier, S; Holowachuk, J; Bonham-Smith, PC; Soroka, J; Lloyd, A, 2006. "HAIRY CANOLA"--*Arabidopsis GL3* induces a dense covering of trichomes on *Brassica napus* seedlings. Plant Mol. Biol. 60:679-698. Pubmed:16649106.
- Gruis, DF; Guo, H; Selinger, D; Tian, Q; Olsen, OA, 2006. Surface position, not signaling from surrounding maternal tissues, specifies aleurone epidermal cell fate in maize. Plant Physiol. 141:898-909. Pubmed:16698897.
- Gu, R; Zhao, L; Zhang, Y; Chen, X; Bao, J; Zhao, J; Wang, Z; Fu, J; Liu, T; Wang, J; Wang, G, 2006. Isolation of a maize beta-glucosidase gene promoter and characterization of its activity in transgenic tobacco. Plant Cell Rep. 25:1157-1165. Pubmed:16770627.
- Guadagnuolo, R; Clegg, J; Ellstrand, NC, 2006. Relative fitness of transgenic vs. non-transgenic maize x teosinte hybrids: a field evaluation. Ecol. Appl. 16:1967-1974. Pubmed:17069387.
- Guerrero-Andrade, O; Loza-Rubio, E; Olivera-Flores, T; Fehervari-Bone, T; Gomez-Lim, MA, 2006. Expression of the Newcastle disease virus fusion protein in transgenic maize and immunological studies. Transgenic Res. 15:455-463. Pubmed:16906446.
- Guillaumie, S; San-Clemente, H; Deswarte, C; Martinez, Y; Lapierre, C; Murigneux, A; Barriere, Y; Pichon, M; Goffner, D, 2007. MAIZEWALL. Database and developmental gene expression profiling of cell wall biosynthesis and assembly in maize. Plant Physiol. 143:339-363. Pubmed:17098859.
- Gunes, A; Inal, A; Alpaslan, M; Eraslan, F; Bagci, EG; Cicek, N, 2006. Salicylic acid induced changes on some physiological parameters symptomatic for oxidative stress and mineral nutrition in maize (*Zea mays* L.) grown under salinity. J. Plant Physiol. (in press). Pubmed:16690163.
- Guo, L; Wang, ZY; Lin, H; Cui, WE; Chen, J; Liu, M; Chen, ZL; Qu, LJ; Gu, H, 2006. Expression and functional analysis of the rice plasma-membrane intrinsic protein gene family. Cell Res. 16:277-286. Pubmed:16541126.
- Guo, M; Rupe, MA; Yang, X; Crasta, O; Zinselmeier, C; Smith, OS; Bowen, B, 2006. Genome-wide transcript analysis of maize hybrids: allelic additive gene expression and yield heterosis. Theor. Appl. Genet. 113:831-845. Pubmed:16868764.
- Guo, SJ; Li, JR; Qiao, WH; Zhang, XS, 2006. Analysis of amylose accumulation during seed development in maize. Yi Chuan Xue Bao 33:1014-1019. Pubmed:17112973.
- Guo, Y; Ma, Y; Zhan, Z; Li, B; Dingkuhn, M; Luquet, D; De Reffye, P, 2006. Parameter optimization and field validation of the functional-structural model GREENLAB for maize. Ann. Bot. (Lond.) 97:217-230. Pubmed:16390847.
- Gupta, S; Ciungu, A; Jameson, N; Lal, SK, 2006. Alternative splicing expression of U1 snRNP 70K gene is evolutionary conserved between different plant species. DNA Seq. 17:254-261. Pubmed:17312944.
- Gustafson, DI; Brants, IO; Horak, MJ; Remund, KM; Rosenbaum, EW; Soteres, JK, 2006. Empirical modeling of genetically modified maize grain production practices to achieve European Union labeling thresholds. Crop Sci. 46:2133-2140. Pubmed:20061219.
- Gutierrez-Marcos, JF; Costa, LM; Evans, MM, 2006. Maternal gametophytic *baseless1* is required for development of the central cell and early endosperm patterning in maize (*Zea mays*). Genetics 174:317-329. Pubmed:16849604.
- Hachez, C; Moshelion, M; Zelazny, E; Cavez, D; Chaumont, F, 2006. Localization and quantification of plasma membrane aquaporin expression in maize primary root: a clue to understanding their role as cellular plumbbers. Plant Mol. Biol. 62:305-323. Pubmed:16845476.
- Hallauer, AR; Smith, JSC, 2006. Dedication: Donald N. Duvick - Maize breeder, geneticist, and administrator. Maydica 51:171-176. Pubmed:20061003.
- Hamant, O; Ma, H; Cande, WZ, 2006. Genetics of meiotic prophase I in plants. Annu. Rev. Plant Biol. 57:267-302. Pubmed:16669763.
- Hamberg, M; Chechetkin, IR; Grechkin, AN; Ponce de Leon, I; Castresana, C; Bannenberg, G, 2006. Synthesis of 3-oxalinolenic acid and beta-oxidation-resistant 3-oxa-oxylipins. Lipids 41:499-506. Pubmed:16933794.
- Han, F; Lamb, JC; Birchler, JA, 2006. High frequency of centromere inactivation resulting in stable dicentric chromosomes of maize. Proc. Natl. Acad. Sci. USA 103:3238-3243. Pubmed:16492777.

- Hanisch, J; Waltermann, M; Robenek, H; Steinbuchel, A, 2006. Eukaryotic lipid body proteins in oleogenous actinomycetes and their targeting to intracellular triacylglycerol inclusions: Impact on models of lipid body biogenesis. *Appl. Environ. Microbiol.* 72:6743-6750. Pubmed:17021226.
- Hardin, SC; Duncan, KA; Huber, SC, 2006. Determination of structural requirements and probable regulatory effectors for membrane association of maize sucrose synthase 1. *Plant Physiol.* 141:1106-1119. Pubmed:16698903.
- Harfouche, AL; Shivaji, R; Stocker, R; Williams, PW; Luthe, DS, 2006. Ethylene signaling mediates a maize defense response to insect herbivory. *Mol. Plant Microbe Interact.* 19:189-199. Pubmed:16529381.
- Haun, WJ; Laoueille-Duprat, S; O'Connell, M, J; Spillane, C; Grossniklaus, U; Phillips, AR; Kaepller, SM; Springer, NM, 2007. Genomic imprinting, methylation and molecular evolution of maize *Enhancer of zeste* (*Mez*) homologs. *Plant J.* 49:325-337. Pubmed:17181776.
- Hawbaker, MS; Goodman, MM, 2006. Resistance of temperately adapted tropical inbred lines and testcrosses to three important maize pathogens. *Maydica* 51:135-139. Pubmed:20061017.
- Hawkins, JS; Kim, H; Nason, JD; Wing, RA; Wendel, JF, 2006. Differential lineage-specific amplification of transposable elements is responsible for genome size variation in *Gossypium*. *Genome Res.* 16:1252-1261. Pubmed:16954538.
- Hayes, ML; Reed, ML; Hegeman, CE; Hanson, MR, 2006. Sequence elements critical for efficient RNA editing of a tobacco chloroplast transcript in vivo and in vitro. *Nucl. Acids Res.* 34:3742-3754. Pubmed:16893957.
- Henderson, DC; Zhang, X; Brooks, L, 3rd; Scanlon, MJ, 2006. *RAGGED SEEDLING2* is required for expression of *KANADI2* and *REVOLUTA* homologues in the maize shoot apex. *Genesis* 44:372-382. Pubmed:16858691.
- Herrmann, MM; Pinto, S; Kluth, J; Wienand, U; Lorbiecke, R, 2006. The PT11-like kinase ZmPt1a from maize (*Zea mays* L.) co-localizes with callose at the plasma membrane of pollen and facilitates a competitive advantage to the male gametophyte. *BMC Plant Biol.* 6:22. Pubmed:17022830.
- Hoecker, N; Keller, B; Piepho, HP; Hochholdinger, F, 2006. Manifestation of heterosis during early maize (*Zea mays* L.) root development. *Theor. Appl. Genet.* 112:421-429. Pubmed:16362278.
- Hola, D; Kocova, M; Rothova, O; Wilhelmova, N; Benesova, M, 2006. Recovery of maize (*Zea mays* L.) inbreds and hybrids from chilling stress of various duration: Photosynthesis and antioxidant enzymes. *J. Plant Physiol.* (in press). Pubmed:16884820.
- Holding, DR; Larkins, BA, 2006. The development and importance of zein protein bodies in maize endosperm. *Maydica* 51:243-254. Pubmed:20061003.
- Holland, JB; Bretting, PK; Bubeck, DM; Cardinal, AJ; Holley, RN; Uhr, DV, 2006. Major M. Goodman - A laudation. *Maydica* 51:3-13. Pubmed:20061017.
- Hu, Q; Nelson, K; Luo, H, 2006. FLP-mediated site-specific recombination for genome modification in turfgrass. *Biotechnol. Lett.* 28:1793-1804. Pubmed:16912917.
- Hu, X; Zhang, A; Zhang, J; Jiang, M, 2006. Abscisic acid is a key inducer of hydrogen peroxide production in leaves of maize plants exposed to water stress. *Plant Cell Physiol.* 47:1484-1495. Pubmed:16990290.
- Hu, YM; Tang, JH; Yang, H; Xie, HL; Lu, XM; Niu, JH; Chen, WC, 2006. Identification and mapping of *Rf-I* an inhibitor of the *Rf5* restorer gene for *Cms-C* in maize (*Zea mays* L.). *Theor. Appl. Genet.* 113:357-360. Pubmed:16791701.
- Huang, S; Frizzi, A; Florida, CA; Kruger, DE; Luethy, MH, 2006. High lysine and high tryptophan transgenic maize resulting from the reduction of both 19- and 22-kD alpha-zeins. *Plant Mol. Biol.* 61:525-535. Pubmed:16830184.
- Huo, N; Gu, YQ; Lazo, GR; Vogel, JP; Coleman-Derr, D; Luo, MC; Thilmony, R; Garvin, DF; Anderson, OD, 2006. Construction and characterization of two BAC libraries from *Brachypodium distachyon*, a new model for grass genomics. *Genome* 49:1099-1108. Pubmed:17110990.
- Ilic, K; Kellogg, EA; Jaiswal, P; Zapata, F; Stevens, PF; Vincent, LP; Avraham, S; Reiser, L; Pujar, A; Sachs, MM; Whitman, NT; McCouch, SR; Schaeffer, ML; Ware, DH; Stein, LD; Rhee, SY, 2007. The plant structure ontology, a unified vocabulary of anatomy and morphology of a flowering plant. *Plant Physiol.* 143:587-599. Pubmed:17142475.
- Irish, EE; McMurray, D, 2006. Rejuvenation by shoot apex culture recapitulates the developmental increase of methylation at the maize gene *Pl-Blotched*. *Plant Mol. Biol.* 60:747-758. Pubmed:16649110.
- Jabeen, R; Yamada, K; Shigemori, H; Hasegawa, T; Hara, M; Kuboi, T; Hasegawa, K, 2006. Induction of beta-glucosidase activity in maize coleoptiles by blue light illumination. *J. Plant Physiol.* 163:538-545. Pubmed:16473658.
- Jaiswal, P; Ni, J; Yap, I; Ware, D; Spooner, W; Youens-Clarkson, K; Ren, L; Liang, C; Zhao, W; Ratnapu, K; Faga, B; Canaran, P; Fogleman, M; Hebbard, C; Avraham, S; Schmidt, S; Casstevens, TM; Buckler, ES; Stein, L; McCouch, S, 2006. Gramene: a bird's eye view of cereal genomes. *Nucl. Acids Res.* 34:D717-723. Pubmed:16381966.
- Jia, J; Fu, J; Zheng, J; Zhou, X; Huai, J; Wang, J; Wang, M; Zhang, Y; Chen, X; Zhang, J; Zhao, J; Su, Z; Lv, Y; Wang, G, 2006. Annotation and expression profile analysis of 2073 full-length cDNAs from stress-induced maize (*Zea mays* L.) seedlings. *Plant J.* 48:710-727. Pubmed:17076806.
- Jiang, K; Ballinger, T; Li, D; Zhang, S; Feldman, L, 2006. A role for mitochondria in the establishment and maintenance of the maize root quiescent center. *Plant Physiol.* 140:1118-1125. Pubmed:16443698.
- Jiang, K; Zhang, S; Lee, S; Tsai, G; Kim, K; Huang, H; Chilcott, C; Zhu, T; Feldman, LJ, 2006. Transcription profile analyses identify genes and pathways central to root cap functions in maize. *Plant Mol. Biol.* 60:343-363. Pubmed:16514559.
- Jines, MP; Balint-Kurti, P; Robertson-Hoyt, LA; Molnar, T; Holland, JB; Goodman, MM, 2007. Mapping resistance to Southern rust in a tropical by temperate maize recombinant inbred topcross population. *Theor. Appl. Genet.* 114:659-667. Pubmed:17177063.

- Johnson, C; Bowman, L; Adai, AT; Vance, V; Sundaresan, V, 2007. CSRDB: a small RNA integrated database and browser resource for cereals. *Nucl. Acids Res.* 35:D829-833. Pubmed:17169981.
- Jones, DL; Blancaflor, EB; Kochian, LV; Gilroy, S, 2006. Spatial coordination of aluminium uptake, production of reactive oxygen species, callose production and wall rigidification in maize roots. *Plant Cell Environ.* 29:1309-1318. Pubmed:17080952.
- Ju, C; Zhang, F; Gao, Y; Zhang, W; Yan, J; Dai, J; Li, J, 2006. Cloning, chromosome mapping and expression analysis of an *R2R3-MYB* gene under-expressed in maize hybrid. *Mol. Biol. Rep.* 33:103-110. Pubmed:16817019.
- Jung, HG; Casler, MD, 2006. Maize stem tissues: Impact of development on cell wall degradability. *Crop Sci.* 46:1801-1809. Pubmed:20061017.
- Jung, HG; Casler, MD, 2006. Maize stem tissues: Cell wall concentration and composition during development. *Crop Sci.* 46:1793-1800. Pubmed:20061017.
- Kalyanaraman, A; Aluru, S; Schnable, PS, 2006. Turning repeats to advantage: scaffolding genomic contigs using LTR retrotransposons. *Comput. Syst. Bioinformatics Conf.* 167-178. Pubmed:17369635.
- Kamara, AY; Menkir, A; Kureh, I; Omoigui, LO, 2006. Response to low soil nitrogen stress of S-1, maize breeding lines, selected for high vertical root-pulling resistance. *Maydica* 51:425-433. Pubmed:20070123.
- Karkonen, A; Fry, SC, 2006. Novel characteristics of UDP-glucose dehydrogenase activities in maize: non-involvement of alcohol dehydrogenases in cell wall polysaccharide biosynthesis. *Planta* 223:858-870. Pubmed:16453102.
- Kato, A; Albert, PS; Vega, JM; Birchler, JA, 2006. Sensitive fluorescence in situ hybridization signal detection in maize using directly labeled probes produced by high concentration DNA polymerase nick translation. *Biotech. Histochem.* 81:71-78. Pubmed:16908431.
- Kato, A; Birchler, JA, 2006. Induction of tetraploid derivatives of maize inbred lines by nitrous oxide gas treatment. *J. Hered.* 97:39-44. Pubmed:16394254.
- Kawakatsu, T; Itoh, J; Miyoshi, K; Kurata, N; Alvarez, N; Veit, B; Nagato, Y, 2006. *PLASTOCHRON2* regulates leaf initiation and maturation in rice. *Plant Cell* 18:612-625. Pubmed:16461585.
- Kellogg, EA, 2007. Floral displays: genetic control of grass inflorescences. *Curr. Opin. Plant Biol.* 10:26-31. Pubmed:17140843.
- Kermicle, JL, 2006. A selfish gene governing pollen-pistil compatibility confers reproductive isolation between maize relatives. *Genetics* 172:499-506. Pubmed:16157680.
- Kermicle, JL; Taba, S; Evans, MMS, 2006. The *gametophyte-1* locus and reproductive isolation among *Zea mays* subspecies. *Maydica* 51:219-225. Pubmed:20061003.
- Kessler, S; Townsley, B; Sinha, N, 2006. L1 division and differentiation patterns influence shoot apical meristem maintenance. *Plant Physiol.* 141:1349-1362. Pubmed:16798950.
- Kidner, C; Timmermans, M, 2006. In situ hybridization as a tool to study the role of microRNAs in plant development. *Methods Mol. Biol.* 342:159-179. Pubmed:16957374.
- Kim, BG; Lee, Y; Hur, HG; Lim, Y; Ahn, JH, 2006. Flavonoid 3'-O-methyltransferase from rice: cDNA cloning, characterization and functional expression. *Phytochemistry* 67:387-394. Pubmed:16412485.
- Kim, CS; Gibbon, BC; Gillikin, JW; Larkins, BA; Boston, RS; Jung, R, 2006. The maize *Mucronate* mutation is a deletion in the 16-kDa gamma-zein gene that induces the unfolded protein response. *Plant J.* 48:440-451. Pubmed:17010110.
- Kim, YH; Woloshuk, CP; Cho, EH; Bae, JM; Song, YS; Huh, GH, 2007. Cloning and functional expression of the gene encoding an inhibitor against *Aspergillus flavus* alpha-amylase, a novel seed lectin from *Lablab purpureus* (*Dolichos lablab*). *Plant Cell Rep.* 26:395-405. Pubmed:17149640.
- Kiran, NS; Polanska, L; Fohlerova, R; Mazura, P; Valkova, M; Smeral, M; Zouhar, J; Malbeck, J; Dobrev, PI; Machackova, I; Brzobohaty, B, 2006. Ectopic over-expression of the maize beta-glucosidase Zm-p60.1 perturbs cytokinin homeostasis in transgenic tobacco. *J. Exp. Bot.* 57:985-996. Pubmed:16488914.
- Kirst, M; Caldo, R; Casati, P; Tanimoto, G; Walbot, V; Wise, RP; Buckler, ES, 2006. Genetic diversity contribution to errors in short oligonucleotide microarray analysis. *Plant Biotechnol. J.* 4:489-498. Pubmed:17309725.
- Koch, M; Strobel, E; Tebbe, CC; Heritage, J; Breves, G; Huber, K, 2006. Transgenic maize in the presence of ampicillin modifies the metabolic profile and microbial population structure of bovine rumen fluid in vitro. *Br. J. Nutr.* 96:820-829. Pubmed:17092369.
- Kollner, TG; O'Maille, PE; Gatto, N; Boland, W; Gershenson, J; Degenhardt, J, 2006. Two pockets in the active site of maize sesquiterpene synthase TPS4 carry out sequential parts of the reaction scheme resulting in multiple products. *Arch. Biochem. Biophys.* 448:83-92. Pubmed:16297849.
- Koterniak, VV, 2006. [A modifier of the *Bg* element of the *Bg-rbg* transposable element system of maize]. *Genetika* 42:185-191. Pubmed:16583702.
- Kozhukhova, NE; Sivolap lu, M, 2006. [Molecular markers in genetic and selection studies of maize]. *Tsitol. Genet.* 40:69-80. Pubmed:17385419.
- Krakowsky, MD; Lee, M; Coors, JG, 2006. Quantitative trait loci for cell wall components in recombinant inbred lines of maize (*Zea mays* L.) II: leaf sheath tissue. *Theor. Appl. Genet.* 112:717-726. Pubmed:16362276.
- Krakowsky, MD; Lee, M; Garay, L; Woodman-Clikeman, W; Long, MJ; Sharopova, N; Frame, B; Wang, K, 2006. Quantitative trait loci for callus initiation and totipotency in maize (*Zea mays* L.). *Theor. Appl. Genet.* 113:821-830. Pubmed:16896717.
- Kriechbaumer, V; Park, WJ; Gierl, A; Glawischnig, E, 2006. Auxin biosynthesis in maize. *Plant Biol. (Stuttg.)* 8:334-339. Pubmed:16807825.

- Kume, T; Sekiya, N; Yano, K, 2006. Heterogeneity in spatial P-distribution and foraging capability by *Zea mays*: effects of patch size and barriers to restrict root proliferation within a patch. *Ann. Bot. (Lond.)* 98:1271-1277. Pubmed:17008353.
- Kunert, R; Gach, JS; Vorauer-Uhl, K; Engel, E; Katinger, H, 2006. Validated method for quantification of genetically modified organisms in samples of maize flour. *J. Agric. Food Chem.* 54:678-681. Pubmed:16448167.
- Kutschera, U; Niklas, KJ, 2007. Photosynthesis research on yellowtops: Macroevolution in progress. *Theory Biosci.* 125:81-92. Pubmed:17412289.
- La Paz, JL; Garcia-Muniz, N; Nadal, A; Esteve, T; Puigdomenech, P; Pla, M, 2006. Interlaboratory transfer of a real-time polymerase chain reaction assay for quantitative detection of genetically modified maize event TC-1507. *J. AOAC Int.* 89:1347-1352. Pubmed:17042186.
- Lamb, JC; Birchler, JA, 2006. Retroelement genome painting: cytological visualization of retroelement expansions in the genera *Zea* and *Tripsacum*. *Genetics* 173:1007-1021. Pubmed:16582446.
- Landi, P; Sanguineti, MC; Liu, C; Li, Y; Wang, TY; Giuliani, S; Bellotti, M; Salvi, S; Tuberosa, R, 2007. Root-ABA1 QTL affects root lodging, grain yield, and other agronomic traits in maize grown under well-watered and water-stressed conditions. *J. Exp. Bot.* 58:319-326. Pubmed:17050640.
- Law, RD; Russell, DA; Thompson, LC; Schroeder, SC; Middle, CM; Tremaine, MT; Jury, TP; Delannay, X; Slater, SC, 2006. Biochemical limitations to high-level expression of humanized monoclonal antibodies in transgenic maize seed endosperm. *Biochim. Biophys. Acta* 1760:1434-1444. Pubmed:16842925.
- Lawrence, CJ; Seifried, TE; Bass, HW; Anderson, LK, 2006. Predicting chromosomal locations of genetically mapped loci in maize using the Morgan2McClintock Translator. *Genetics* 172:2007-2009. Pubmed:16387866.
- Le Clerc, V; Cadot, V; Canadas, M; Lallemand, J; Guerin, D; Boulineau, F, 2006. Indicators to assess temporal genetic diversity in the French Catalogue: no losses for maize and peas. *Theor. Appl. Genet.* 113:1197-1209. Pubmed:16900350.
- Leakey, AD; Uribelarrea, M; Ainsworth, EA; Naidu, SL; Rogers, A; Ort, DR; Long, SP, 2006. Photosynthesis, productivity, and yield of maize are not affected by open-air elevation of CO<sub>2</sub> concentration in the absence of drought. *Plant Physiol.* 140:779-790. Pubmed:16407441.
- LeDeaux, JR; Graham, GI; Stuber, CW, 2006. Stability of QTLs involved in heterosis in maize when mapped under several stress conditions. *Maydica* 51:151-167. Pubmed:20061017.
- Lee, EA; Chakravarty, R; Good, B; Ash, MJ; Kannenberg, LW, 2006. Registration of 38 maize (*Zea mays* L.) breeding populations adapted to short-season environments. *Crop Sci.* 46:2728-2733. Pubmed:20070306.
- Lee, EA; Singh, A; Ash, MJ; Good, B, 2006. Use of sister-lines and the performance of modified single-cross maize hybrids. *Crop Sci.* 46:312-320. Pubmed:20060524.
- Lee, JK; Park, JY; Kim, JH; Kwon, SJ; Shin, JH; Hong, SK; Min, HK; Kim, NS, 2006. Genetic mapping of the *Isaac*-CACTA transposon in maize. *Theor. Appl. Genet.* 113:16-22. Pubmed:16783589.
- Lee, KM; Bean, SR; Alavi, S; Herman, TJ; Waniska, RD, 2006. Physical and biochemical properties of maize hardness and extrudates of selected hybrids. *J. Agric. Food Chem.* 54:4260-4269. Pubmed:16756355.
- Lee, SH; Hamaker, BR, 2006. Cys155 of 27 kDa maize gamma-zein is a key amino acid to improve its in vitro digestibility. *FEBS Lett.* 580:5803-5806. Pubmed:17045266.
- Lee, SH; Min, DM; Kim, JK, 2006. Qualitative and quantitative polymerase chain reaction analysis for genetically modified maize MON863. *J. Agric. Food Chem.* 54:1124-1129. Pubmed:16478226.
- Lee, YH; Tamura, K; Maeda, M; Hoshino, M; Sakurai, K; Takahashi, S; Ikegami, T; Hase, T; Goto, Y, 2007. Cores and pH-dependent dynamics of ferredoxin-NADP<sup>+</sup> reductase revealed by hydrogen/deuterium exchange. *J. Biol. Chem.* 282:5959-5967. Pubmed:17192259.
- Lekacz, H; Karcz, W, 2006. The effect of auxins (IAA and 4-Cl-IAA) on the redox activity and medium pH of *Zea mays* L. root segments. *Cell. Mol. Biol. Lett.* 11:376-383. Pubmed:16847555.
- Li, K; Yang, J; Liu, J; Du, X; Wei, C; Su, W; He, G; Zhang, Q; Hong, F; Qian, X, 2006. Cloning, characterization and tissue-specific expression of a cDNA encoding a novel *EMBRYONIC FLOWER* 2 gene (*OsEMF2*) in *Oryza sativa*. *DNA Seq.* 17:74-78. Pubmed:16753820.
- Li, MS; Li, XH; Salvi, S; Tuberosa, R; Yuan, LX; Rotondo, F; Bai, L; Zhang, SH, 2006. Genetic relationships among CIMMYT subtropical QPM and Chinese maize inbred lines based on SSRs. *Maydica* 51:543-549. Pubmed:20070123.
- Li, W; Gill, BS, 2006. Multiple genetic pathways for seed shattering in the grasses. *Funct. Integr. Genomics* 6:300-309. Pubmed:16404644.
- Li, W; Ruf, S; Bock, R, 2006. Constancy of organellar genome copy numbers during leaf development and senescence in higher plants. *Mol. Genet. Genomics* 275:185-192. Pubmed:16308694.
- Li, XP; Gan, R; Li, PL; Ma, YY; Zhang, LW; Zhang, R; Wang, Y; Wang, NN, 2006. Identification and functional characterization of a leucine-rich repeat receptor-like kinase gene that is involved in regulation of soybean leaf senescence. *Plant Mol. Biol.* 61:829-844. Pubmed:16927199.
- Liang, M; Haroldsen, V; Cai, X; Wu, Y, 2006. Expression of a putative laccase gene, *ZmLAC1*, in maize primary roots under stress. *Plant Cell Environ.* 29:746-753. Pubmed:17087459.

- Libourel, IG; van Bodegom, PM; Fricker, MD; Ratcliffe, RG, 2006. Nitrite reduces cytoplasmic acidosis under anoxia. *Plant Physiol.* 142:1710-1717. Pubmed:17071644.
- Lim, TS; Chitra, TR; Han, P; Pua, EC; Yu, H, 2006. Cloning and characterization of *Arabidopsis* and *Brassica juncea* flavin-containing amine oxidases. *J. Exp. Bot.* 57:4155-4169. Pubmed:17122409.
- Lincoln, T, 2006. Chemical ecology: in defence of maize. *Nature* 439:278. Pubmed:16421557.
- Lipsius, K; Wilhelm, R; Richter, O; Schmalstieg, KJ; Schiemann, J, 2006. Meteorological input data requirements to predict cross-pollination of GMO Maize with Lagrangian approaches. *Environ. Biosafety Res.* 5:151-168. Pubmed:17445511.
- Liu, Y; Lamkemeyer, T; Jakob, A; Mi, G; Zhang, F; Nordheim, A; Hochholdinger, F, 2006. Comparative proteome analyses of maize (*Zea mays* L.) primary roots prior to lateral root initiation reveal differential protein expression in the lateral root initiation mutant *rum1*. *Proteomics* 6:4300-4308. Pubmed:16819721.
- Longin, CF; Utz, HF; Melchinger, AE; Reif, JC, 2007. Hybrid maize breeding with doubled haploids: II. Optimum type and number of testers in two-stage selection for general combining ability. *Theor. Appl. Genet.* 114:393-402. Pubmed:17180379.
- Longin, CF; Utz, HF; Reif, JC; Schipprack, W; Melchinger, AE, 2006. Hybrid maize breeding with doubled haploids: I. One-stage versus two-stage selection for testcross performance. *Theor. Appl. Genet.* 112:903-912. Pubmed:16435127.
- Loureiro, J; Rodriguez, E; Dolezel, J; Santos, C, 2006. Flow cytometric and microscopic analysis of the effect of tannic acid on plant nuclei and estimation of DNA content. *Ann. Bot. (Lond.)* 98:515-527. Pubmed:16820406.
- Lozovaya, V; Ulanov, A; Lygin, A; Duncan, D; Widholm, J, 2006. Biochemical features of maize tissues with different capacities to regenerate plants. *Planta* 224:1385-1399. Pubmed:16941117.
- Lu, YQ; Ye, ZH; Wu, WR, 2006. Analysis of the phylogenetic relationships among several species of Gramineae using ACGM markers. *Yi Chuan Xue Bao* 33:1127-1131. Pubmed:17185173.
- Luce, AC; Sharma, A; Mollere, OS; Wolfgruber, TK; Nagaki, K; Jiang, J; Presting, GG; Dawe, RK, 2006. Precise centromere mapping using a combination of repeat junction markers and chromatin immunoprecipitation-polymerase chain reaction. *Genetics* 174:1057-1061. Pubmed:16951073.
- Luo, K; Deng, W; Xiao, Y; Zheng, X; Li, Y; Pei, Y, 2006. Leaf senescence is delayed in tobacco plants expressing the maize *knotted1* gene under the control of a wound-inducible promoter. *Plant Cell Rep.* 25:1246-1254. Pubmed:16794826.
- Luo, K; Zheng, X; Chen, Y; Xiao, Y; Zhao, D; McAvoy, R; Pei, Y; Li, Y, 2006. The maize *Knotted1* gene is an effective positive selectable marker gene for *Agrobacterium*-mediated tobacco transformation. *Plant Cell Rep.* 25:403-409. Pubmed:16369767.
- Lutfiyya, LL; Xu, N; D'Ordine, RL; Morrell, JA; Miller, PW; Duff, SM, 2006. Phylogenetic and expression analysis of sucrose phosphate synthase isozymes in plants. *J. Plant Physiol.* (in press). Pubmed:16876912.
- Lutz, B; Wiedemann, S; Albrecht, C, 2006. Degradation of transgenic Cry1Ab DNA and protein in Bt-176 maize during the ensiling process. *J. Anim. Physiol. Anim. Nutr. (Berl.)* 90:116-123. Pubmed:16519756.
- Ma, J; Morrow, DJ; Fernandes, J; Walbot, V, 2006. Comparative profiling of the sense and antisense transcriptome of maize lines. *Genome Biol.* 7:R22. Pubmed:16542496.
- Ma, Y; Li, B; Zhan, Z; Guo, Y; Luquet, D; de Reffye, P; Dingkuhn, M, 2007. Parameter stability of the functional-structural plant model GREENLAB as affected by variation within populations, among seasons and among growth stages. *Ann. Bot. (Lond.)* 99:61-73. Pubmed:17158141.
- Malcomber, ST; Kellogg, EA, 2006. Evolution of unisexual flowers in grasses (Poaceae) and the putative sex-determination gene, *TASSELSEED2 (TS2)*. *New Phytol.* 170:885-899. Pubmed:16684246.
- Malysheva-Otto, LV; Ganal, MW; Roder, MS, 2006. Analysis of molecular diversity, population structure and linkage disequilibrium in a worldwide survey of cultivated barley germplasm (*Hordeum vulgare* L.). *BMC Genet.* 7:6. Pubmed:16433922.
- Manetti, C; Bianchetti, C; Casciani, L; Castro, C; Di Cocco, ME; Miccheli, A; Motto, M; Conti, F, 2006. A metabonomic study of transgenic maize (*Zea mays*) seeds revealed variations in osmolytes and branched amino acids. *J. Exp. Bot.* 57:2613-2625. Pubmed:16831843.
- Maresh, J; Zhang, J; Lynn, DG, 2006. The innate immunity of maize and the dynamic chemical strategies regulating two-component signal transduction in *Agrobacterium tumefaciens*. *ACS Chem. Biol.* 1:165-175. Pubmed:17163664.
- Martin, A; Lee, J; Kichey, T; Gerentes, D; Zivy, M; Tatout, C; Dubois, F; Balliau, T; Valot, B; Davanture, M; Terce-Laforgue, T; Quillere, I; Coque, M; Gallais, A; Gonzalez-Moro, MB; Bethencourt, L; Habash, DZ; Lea, PJ; Charcosset, A; Perez, P; Murigneux, A; Sakakibara, H; Edwards, KJ; Hirel, B, 2006. Two cytosolic glutamine synthetase isoforms of maize are specifically involved in the control of grain production. *Plant Cell* 18:3252-3274. Pubmed:17138698.
- Masi, A; Destro, T; Turetta, L; Varotto, S; Caporale, G; Ferretti, M, 2006. Localization of gamma-glutamyl transferase activity and protein in *Zea mays* organs and tissues. *J. Plant Physiol.* (in press). Pubmed:17074415.
- Maurer, HP; Knaak, C; Melchinger, AE; Ouzunova, M; Frisch, M, 2006. Linkage disequilibrium between SSR markers in six pools of elite lines of an European breeding program for hybrid maize. *Maydica* 51:269-279. Pubmed:20061003.
- McCartney, L; Blake, AW; Flint, J; Bolam, DN; Boraston, AB; Gilbert, HJ; Knox, JP, 2006. Differential recognition of plant cell walls by microbial xylan-specific carbohydrate-binding modules. *Proc. Natl. Acad. Sci. USA* 103:4765-4770. Pubmed:16537424.
- McGinnis, KM; Springer, C; Lin, Y; Carey, CC; Chandler, V, 2006. Transcriptionally silenced transgenes in maize are activated by three mutations defective in paramutation. *Genetics* 173:1637-1647. Pubmed:16702420.
- Menkir, A; Kling, JG; Badu-Apraku, B; Ibikunle, O, 2006. Registration of 26 tropical maize germplasm lines with resistance to *Striga hermonthica*. *Crop Sci.* 46:1007-1009. Pubmed:20060718.

- Menkir, A; Olowolafe, MO; Ingelbrecht, I; Fawole, I; Badu-Apraku, B; Vroh, BI, 2006. Assessment of testcross performance and genetic diversity of yellow endosperm maize lines derived from adapted x exotic backcrosses. *Theor. Appl. Genet.* 113:90-99. Pubmed:16614832.
- Meseka, SK; Menkir, A; Ibrahim, AES; Ajala, SO, 2006. Genetic analysis of performance of maize inbred lines selected for tolerance to drought under low nitrogen. *Maydica* 51:487-495. Pubmed:20070123.
- Messeguer, J; Penas, G; Ballester, J; Bas, M; Serra, J; Salvia, J; Palaudelmas, M; Mele, E, 2006. Pollen-mediated gene flow in maize in real situations of coexistence. *Plant Biotechnol. J.* 4:633-645. Pubmed:17309734.
- Messing, J; Dooner, HK, 2006. Organization and variability of the maize genome. *Curr. Opin. Plant Biol.* 9:157-163. Pubmed:16459130.
- Meyer, S; Pospisil, H; Scholten, S, 2007. Heterosis associated gene expression in maize embryos 6 days after fertilization exhibits additive, dominant and overdominant pattern. *Plant Mol. Biol.* 63:381-391. Pubmed:17082872.
- Mezard, C, 2006. Meiotic recombination hotspots in plants. *Biochem. Soc. Trans.* 34:531-534. Pubmed:16856852.
- Mica, E; Gianfranceschi, L; Pe, ME, 2006. Characterization of five microRNA families in maize. *J. Exp. Bot.* 57:2601-2612. Pubmed:16820394.
- Mohan, S; Ma, PW; Pechan, T; Bassford, ER; Williams, WP; Luthe, DS, 2006. Degradation of the *S. frugiperda* peritrophic matrix by an inducible maize cysteine protease. *J. Insect Physiol.* 52:21-28. Pubmed:16243350.
- Momcilovic, I; Ristic, Z, 2007. Expression of chloroplast protein synthesis elongation factor, EF-Tu, in two lines of maize with contrasting tolerance to heat stress during early stages of plant development. *J. Plant Physiol.* 164:90-99. Pubmed:16542752.
- Monjardino, P; Smith, A-G; Jones, R-J, 2006. Zein transcription and endoreduplication in maize endosperm are differentially affected by heat stress. *Crop Sci.* 46:2581-2589. Pubmed:20070306.
- Monma, K; Moriuchi, R; Sagi, N; Ichikawa, H; Satoh, K; Tobe, T; Kamata, K, 2006. Investigation of false-positive reactions for CBH351 maize in screening PCR analysis. *Shokuhin Eiseigaku Zasshi* 47:9-14. Pubmed:16619851.
- Monneveux, P; Sanchez, C; Beck, D; Edmeades, GO, 2006. Drought tolerance improvement in tropical maize source populations: Evidence of progress. *Crop Sci.* 46:180-191. Pubmed:20060524.
- Morrell, PL; Toleno, DM; Lundy, KE; Clegg, MT, 2006. Estimating the contribution of mutation, recombination and gene conversion in the generation of haplotypic diversity. *Genetics* 173:1705-1723. Pubmed:16624913.
- Morton, BR; Bi, IV; McMullen, MD; Gaut, BS, 2006. Variation in mutation dynamics across the maize genome as a function of regional and flanking base composition. *Genetics* 172:569-577. Pubmed:16219784.
- Mroczeck, RJ; Melo, JR; Luce, AC; Hiatt, EN; Dawe, RK, 2006. The maize Ab10 meiotic drive system maps to supernumerary sequences in a large complex haplotype. *Genetics* 174:145-154. Pubmed:16849609.
- Muehlbauer, GJ; Bhau, BS; Syed, NH; Heinen, S; Cho, S; Marshall, D; Pateyron, S; Buisine, N; Chalhoub, B; Flavell, AJ, 2006. A hAT superfamily transposase recruited by the cereal grass genome. *Mol. Genet. Genomics* 275:553-563. Pubmed:16468023.
- Mulder, C; Wouterse, M; Raubuch, M; Roelofs, W; Rutgers, M, 2006. Can transgenic maize affect soil microbial communities? *PLoS Comput. Biol.* 2:e128. Pubmed:17009863.
- Muller, B; Bourdais, G; Reidy, B; Bencivenni, C; Massonneau, A; Condamine, P; Rolland, G; Conejero, G; Rogowsky, P; Tardieu, F, 2007. Association of specific expansins with growth in maize leaves is maintained under environmental, genetic, and developmental sources of variation. *Plant Physiol.* 143:278-290. Pubmed:17098857.
- Muller, S; Han, S; Smith, LG, 2006. Two kinesins are involved in the spatial control of cytokinesis in *Arabidopsis thaliana*. *Curr. Biol.* 16:888-894. Pubmed:16682350.
- Mulvenna, JP; Mylne, JS; Bharathi, R; Burton, RA; Shirley, NJ; Fincher, GB; Anderson, MA; Craik, DJ, 2006. Discovery of cyclotide-like protein sequences in graminaceous crop plants: ancestral precursors of circular proteins? *Plant Cell* 18:2134-2144. Pubmed:16935986.
- Muniz, LM; Royo, J; Gomez, E; Barrero, C; Bergareche, D; Hueros, G, 2006. The maize transfer cell-specific type-A response regulator *ZmTCRR-1* appears to be involved in intercellular signalling. *Plant J.* 48:17-27. Pubmed:16925601.
- Murata, Y; Ma, JF; Yamaji, N; Ueno, D; Nomoto, K; Iwashita, T, 2006. A specific transporter for iron(III)-phytosiderophore in barley roots. *Plant J.* 46:563-572. Pubmed:16640594.
- Musser, FR; Nyrop, JP; Shelton, AM, 2006. Integrating biological and chemical controls in decision making: European corn borer (Lepidoptera: Crambidae) control in sweet corn as an example. *J. Econ. Entomol.* 99:1538-1549. Pubmed:17066781.
- Muszynski, MG; Dam, T; Li, B; Shirbroun, DM; Hou, Z; Bruggemann, E; Archibald, R; Ananiev, EV; Danilevskaya, ON, 2006. *delayed flowering1* encodes a basic leucine zipper protein that mediates floral inductive signals at the shoot apex in maize. *Plant Physiol.* 142:1523-1536. Pubmed:17071646.
- Nadal, A; Coll, A; La Paz, JL; Esteve, T; Pla, M, 2006. A new PCR-CGE (size and color) method for simultaneous detection of genetically modified maize events. *Electrophoresis* 27:3879-3888. Pubmed:16972302.
- Naef, A; Zesiger, T; Defago, G, 2006. Impact of transgenic Bt maize residues on the mycotoxicogenic plant pathogen *Fusarium graminearum* and the biocontrol agent *Trichoderma atroviride*. *J. Environ. Qual.* 35:1001-1009. Pubmed:16738384.
- Nagy, R; Vasconcelos, MJ; Zhao, S; McElver, J; Bruce, W; Amrhein, N; Raghothama, KG; Bucher, M, 2006. Differential regulation of five Pht1 phosphate transporters from maize (*Zea mays* L.). *Plant Biol. (Stuttg.)* 8:186-197. Pubmed:16547863.
- Nardmann, J; Werr, W, 2006. The shoot stem cell niche in angiosperms: expression patterns of *WUS* orthologues in rice and maize imply major modifications in the course of mono- and dicot evolution. *Mol. Biol. Evol.* 23:2492-2504. Pubmed:16987950.

- Nelson, PT; Jines, MP; Goodman, MM, 2006. Selecting among available, elite tropical maize inbreds for use in long-term temperate breeding. *Maydica* 51:255-262. Pubmed:20061003.
- Nemat Alla, MM; Hassan, NM, 2006. Changes of antioxidants levels in two maize lines following atrazine treatments. *Plant Physiol. Biochem.* 44:202-210. Pubmed:16777423.
- Nemchenko, A; Kunze, S; Feussner, I; Kolomiets, M, 2006. Duplicate maize 13-lipoxygenase genes are differentially regulated by circadian rhythm, cold stress, wounding, pathogen infection, and hormonal treatments. *J. Exp. Bot.* 57:3767-3779. Pubmed:17005920.
- Nishimaru, T, 2006. [Study on the structure activity relationship of a phytosiderophore, mugineic acid]. *Yakugaku Zasshi* 126:473-479. Pubmed:16819268.
- Nishimura, T; Mori, Y; Furukawa, T; Kadota, A; Koshiba, T, 2006. Red light causes a reduction in IAA levels at the apical tip by inhibiting de novo biosynthesis from tryptophan in maize coleoptiles. *Planta* 224:1427-1435. Pubmed:16741747.
- Niu, J; Chen, F; Mi, G; Li, C; Zhang, F, 2007. Transpiration, and nitrogen uptake and flow in two maize (*Zea mays* L.) inbred lines as affected by nitrogen supply. *Ann. Bot. (Lond.)* 99:153-160. Pubmed:17088295.
- Nocito, FF; Lancilli, C; Crema, B; Fourcroy, P; Davidian, JC; Sacchi, GA, 2006. Heavy metal stress and sulfate uptake in maize roots. *Plant Physiol.* 141:1138-1148. Pubmed:16698905.
- Nogueira, FT; Sarkar, AK; Chitwood, DH; Timmermans, MC, 2006. Organ polarity in plants Is specified through the opposing activity of two distinct small regulatory RNAs. *Cold Spring Harbor Symp. Quant. Biol.* 71:157-164. Pubmed:17381292.
- Obrist, LB; Klein, H; Dutton, A; Bigler, F, 2006. Assessing the effects of Bt Maize on the predatory mite *Neoseiulus cucumeris*. *Exp. Appl. Acarol.* 38:125-139. Pubmed:16596347.
- Odland, W; Baumgarten, A; Phillips, R, 2006. Ancestral rice blocks define multiple related regions in the maize genome. *Crop Sci.* 46:S41-S48. Pubmed:20070306.
- Offermann, S; Danker, T; Dreymuller, D; Kalamajka, R; Topsch, S; Weyand, K; Peterhansel, C, 2006. Illumination is necessary and sufficient to induce histone acetylation independent of transcriptional activity at the C4-specific phosphoenolpyruvate carboxylase promoter in maize. *Plant Physiol.* 141:1078-1088. Pubmed:16679423.
- Ogo, Y; Itai, RN; Nakanishi, H; Inoue, H; Kobayashi, T; Suzuki, M; Takahashi, M; Mori, S; Nishizawa, NK, 2006. Isolation and characterization of IRO2, a novel iron-regulated bHLH transcription factor in graminaceous plants. *J. Exp. Bot.* 57:2867-2878. Pubmed:16887895.
- Oide, S; Moeder, W; Krasnoff, S; Gibson, D; Haas, H; Yoshioka, K; Turgeon, BG, 2006. NPS6, encoding a nonribosomal peptide synthetase involved in siderophore-mediated iron metabolism, is a conserved virulence determinant of plant pathogenic ascomycetes. *Plant Cell* 18:2836-2853. Pubmed:17056706.
- Okada, T; Bhalla, PL; Singh, MB, 2006. Expressed sequence tag analysis of *Lilium longiflorum* generative cells. *Plant Cell Physiol.* 47:698-705. Pubmed:16571618.
- Oldenburg, DJ; Rowan, BA; Zhao, L; Walcher, CL; Schleh, M; Bendich, AJ, 2006. Loss or retention of chloroplast DNA in maize seedlings is affected by both light and genotype. *Planta* 225:41-55. Pubmed:16941116.
- Olsen, KM; Caicedo, AL; Polato, N; McClung, A; McCouch, S; Purugganan, MD, 2006. Selection under domestication: evidence for a sweep in the rice waxy genomic region. *Genetics* 173:975-983. Pubmed:16547098.
- Omoigui, LO; Alabi, SO; Ado, SG; Ajala, SO; Kamara, AY, 2006. Genetic gains from cycles of full-sib recurrent selection for low nitrogen tolerance in a tropical maize population. *Maydica* 51:497-505. Pubmed:20070123.
- Ong, HC; Palmer, JD, 2006. Pervasive survival of expressed mitochondrial *rps14* pseudogenes in grasses and their relatives for 80 million years following three functional transfers to the nucleus. *BMC Evol Biol.* 6:55. Pubmed:16842621.
- Ordas, B; Padilla, G; Malvar, RA; Ordas, A; Rodriguez, VM; Revilla, P, 2006. Cold tolerance improvement of sugary *enhancer1* hybrids of sweet corn. *Maydica* 51:567-574. Pubmed:20070123.
- Osmont, KS; Sadeghian, N; Freeling, M, 2006. Mosaic analysis of *extended auricle1* (*eta1*) suggests that a two-way signaling pathway is involved in positioning the blade/sheath boundary in *Zea mays*. *Dev. Biol.* 295:1-12. Pubmed:16684518.
- Ostheimer, GJ; Rojas, M; Hadjivassiliou, H; Barkan, A, 2006. Formation of the CRS2-CAF2 group II intron splicing complex is mediated by a 22-amino acid motif in the COOH-terminal region of CAF2. *J. Biol. Chem.* 281:4732-4738. Pubmed:16379013.
- Ota, H; Lim, TK; Tanaka, T; Yoshino, T; Harada, M; Matsunaga, T, 2006. Automated DNA extraction from genetically modified maize using aminosilane-modified bacterial magnetic particles. *J. Biotechnol.* 125:361-368. Pubmed:16621089.
- Palanivelu, R; Preuss, D, 2006. Distinct short-range ovule signals attract or repel *Arabidopsis thaliana* pollen tubes in vitro. *BMC Plant Biol.* 6:7. Pubmed:16595022.
- Panteris, E; Apostolakos, P; Galatis, B, 2006. Cytoskeletal asymmetry in *Zea mays* subsidiary cell mother cells: a monopolar prophase microtubule half-spindle anchors the nucleus to its polar position. *Cell Motil. Cytoskeleton* 63:696-709. Pubmed:16986138.
- Parasharami, VA; Naik, VB; von Arnold, S; Nadgauwa, RS; Clapham, DH, 2006. Stable transformation of mature zygotic embryos and regeneration of transgenic plants of chir pine (*Pinus roxburghii* Sarg.). *Plant Cell Rep.* 24:708-714. Pubmed:16133348.
- Parida, SK; Anand Raj Kumar, K; Dalal, V; Singh, NK; Mohapatra, T, 2006. Unigene derived microsatellite markers for the cereal genomes. *Theor. Appl. Genet.* 112:808-817. Pubmed:16429310.

- Park, SJ; Piao, HL; Xuan, YH; Park, SH; Je, BI; Kim, CM; Lee, EJ; Ryu, B; Lee, KH; Lee, GH; Nam, MH; Yeo, US; Lee, MC; Yun, DW; Eun, MY; Han, CD, 2006. Analysis of intragenic *Ds* transpositions and excision events generating novel allelic variation in rice. *Mol. Cells* 21:284-293. Pubmed:16682825.
- Parys, E; Jastrzebski, H, 2006. Light-enhanced dark respiration in leaves, isolated cells and protoplasts of various types of C4 plants. *J. Plant Physiol.* 163:638-647. Pubmed:16545997.
- Paszkowski, U; Jakovleva, L; Boller, T, 2006. Maize mutants affected at distinct stages of the arbuscular mycorrhizal symbiosis. *Plant J.* 47:165-173. Pubmed:16762030.
- Pedreschi, R; Cisneros-Zevallos, L, 2006. Antimutagenic and antioxidant properties of phenolic fractions from Andean purple corn (*Zea mays* L.). *J. Agric. Food Chem.* 54:4557-4567. Pubmed:16786998.
- Pereira, LK; Mangolin, CA; Scapim, CA; Pacheco, CAP; Bonato, CM; Machado, MFPS, 2006. Malate dehydrogenase isozyme patterns in four cycles of half-sib selection from CMS-43 popcorn variety (*Zea mays* L.). *Maydica* 51:561-566. Pubmed:20070123.
- Peremyslova, EE, 2006. [Use of homeotic mutation *tasselseed2* for investigation of the action of maize meiotic genes during micro- and megasporogenesis]. *Genetika* 42:519-529. Pubmed:16756071.
- Perin, L; Martinez-Aguilar, L; Castro-Gonzalez, R; Estrada-de Los Santos, P; Cabellos-Avelar, T; Guedes, HV; Reis, VM; Caballero-Mellado, J, 2006. Diazotrophic *Burkholderia* species associated with field-grown maize and sugarcane. *Appl. Environ. Microbiol.* 72:3103-3110. Pubmed:16672447.
- Perry, L; Sandweiss, DH; Piperno, DR; Rademaker, K; Malpass, MA; Umire, A; de la Vera, P, 2006. Early maize agriculture and interzonal interaction in southern Peru. *Nature* 440:76-79. Pubmed:16511492.
- Petersen, A; Dresselhaus, T; Grobe, K; Becker, WM, 2006. Proteome analysis of maize pollen for allergy-relevant components. *Proteomics* 6:6317-6325. Pubmed:17080481.
- Peterson, P-A; Bianchi, A, 2006. Donald N. Duvick: An Illinois farm youth to an academic in the corporate offices. *Maydica* 51:169. Pubmed:20061003.
- Peterson, RK; Meyer, SJ; Wolf, AT; Wolt, JD; Davis, PM, 2006. Genetically engineered plants, endangered species, and risk: a temporal and spatial exposure assessment for Karner blue butterfly larvae and Bt maize pollen. *Risk Anal.* 26:845-858. Pubmed:16834638.
- Phan, BH; Jin, W; Topp, CN; Zhong, CX; Jiang, J; Dawe, RK; Parrott, WA, 2007. Transformation of rice with long DNA-segments consisting of random genomic DNA or centromere-specific DNA. *Transgenic Res.* 16:341-351. Pubmed:17103243.
- Philippar, K; Buchsenschutz, K; Edwards, D; Loffler, J; Luthen, H; Kranz, E; Edwards, KJ; Hedrich, R, 2006. The auxin-induced K(+) channel gene *Zmk1* in maize functions in coleoptile growth and is required for embryo development. *Plant Mol. Biol.* 61:757-768. Pubmed:16897490.
- Piepho, HP; Keller, B; Hoecker, N; Hochholdinger, F, 2006. Combining signals from spotted cDNA microarrays obtained at different scanning intensities. *Bioinformatics* 22:802-807. Pubmed:16418237.
- Pixley, K-V; Dhliwayo, T; Tongona, P, 2006. Improvement of a maize population by full-sib selection alone versus full-sib with selection during inbreeding. *Crop Sci.* 46:1130-1136. Pubmed:20060918.
- Pla, M; La Paz, JL; Penas, G; Garcia, N; Palau-delmas, M; Esteve, T; Messeguer, J; Mele, E, 2006. Assessment of real-time PCR based methods for quantification of pollen-mediated gene flow from GM to conventional maize in a field study. *Transgenic Res.* 15:219-228. Pubmed:16604462.
- Polanska, L; Vicankova, A; Novakova, M; Malbeck, J; Dobrev, PI; Brzobohaty, B; Vankova, R; Machackova, I, 2007. Altered cytokinin metabolism affects cytokinin, auxin, and abscisic acid contents in leaves and chloroplasts, and chloroplast ultrastructure in transgenic tobacco. *J. Exp. Bot.* 58:637-649. Pubmed:17175552.
- Pompa, A; Vitale, A, 2006. Retention of a bean phaseolin/maize gamma-Zein fusion in the endoplasmic reticulum depends on disulfide bond formation. *Plant Cell* 18:2608-2621. Pubmed:17041149.
- Popelkova, H; Fraaije, MW; Novak, O; Frebortova, J; Bilyeu, KD; Frebort, I, 2006. Kinetic and chemical analyses of the cytokinin dehydrogenase-catalysed reaction: correlations with the crystal structure. *Biochem J.* 398:113-124. Pubmed:16686601.
- Porch, TG; Tseung, CW; Schmelz, EA; Mark Settles, A, 2006. The maize *Viviparous10/Viviparous13* locus encodes the *Cnx1* gene required for molybdenum cofactor biosynthesis. *Plant J.* 45:250-263. Pubmed:16367968.
- Poroyko, V; Spollen, WG; Hejlek, LG; Hernandez, AG; LeNoble, ME; Davis, G; Nguyen, HT; Springer, GK; Sharp, RE; Bohnert, HJ, 2007. Comparing regional transcript profiles from maize primary roots under well-watered and low water potential conditions. *J. Exp. Bot.* 58:279-289. Pubmed:16990373.
- Porter, SE; Stoll, DR; Rutan, SC; Carr, PW; Cohen, JD, 2006. Analysis of four-way two-dimensional liquid chromatography-diode array data: Application to metabolomics. *Anal. Chem.* 78:5559-5569. Pubmed:16878896.
- Pratt, RC; Casey, MA, 2006. Registration of maize germplasm line Oh605. *Crop Sci.* 46:1004-1005. Pubmed:20060718.
- Preciado-Ortiz, R; Guerrero, R; Ortega, A; Terron, A; Crossa, F; Cordova, H; Reyes, C; Aguilar, G; Tut, C; Gomez, N; Cervantes, E, 2006. Identification of superior quality protein maize hybrids for different mega-environments using the biplot methodology. *Maydica* 51:451-460. Pubmed:20070123.
- Pret'ova, A; Obert, B; Bartosova, Z, 2006. Haplid formation in maize, barley, flax, and potato. *Protoplasma* 228:107-114. Pubmed:16937062.
- Qi, XM; Li, PJ; Liu, W; Xie, LJ, 2006. Multiple biomarkers response in maize (*Zea mays* L.) during exposure to copper. *J. Environ. Sci. (China)* 18:1182-1188. Pubmed:17294962.

- Rabinowicz, PD; Bennetzen, JL, 2006. The maize genome as a model for efficient sequence analysis of large plant genomes. *Curr. Opin. Plant Biol.* 9:149-156. Pubmed:16459129.
- Rasmussen, CC; Hallauer, AR, 2006. Evaluation of heterotic patterns of Iowa Stiff Stalk Synthetic and non-stiff stalk synthetic maize populations. *Maydica* 51:177-186. Pubmed:20061003.
- Rauschen, S; Schuphan, I, 2006. Fate of the Cry1Ab protein from Bt-maize MON810 silage in biogas production facilities. *J. Agric. Food Chem.* 54:879-883. Pubmed:16448198.
- Reddy, AM; Reddy, VS; Scheffler, BE; Wienand, U; Reddy, AR, 2007. Novel transgenic rice overexpressing anthocyanidin synthase accumulates a mixture of flavonoids leading to an increased antioxidant potential. *Metab. Eng.* 9:95-111. Pubmed:17157544.
- Reif, JC; Warburton, ML; Xia, XC; Hoisington, DA; Crossa, J; Taba, S; Muminovic, J; Bohn, M; Frisch, M; Melchinger, AE, 2006. Grouping of accessions of Mexican races of maize revisited with SSR markers. *Theor. Appl. Genet.* 113:177-185. Pubmed:16791685.
- Remigereau, MS; Robin, O; Siljak-Yakovlev, S; Sarr, A; Robert, T; Langin, T, 2006. *Tuareg*, a novel miniature-inverted repeat family of pearl millet (*Pennisetum glaucum*) related to the *PIF* superfamily of maize. *Genetica* 128:205-216. Pubmed:17028951.
- Ren, H; Gao, Z; Chen, L; Wei, K; Liu, J; Fan, Y; Davies, WJ; Jia, W; Zhang, J, 2007. Dynamic analysis of ABA accumulation in relation to the rate of ABA catabolism in maize tissues under water deficit. *J. Exp. Bot.* 58:211-219. Pubmed:16982652.
- Requejo, R; Tena, M, 2006. Maize response to acute arsenic toxicity as revealed by proteome analysis of plant shoots. *Proteomics* 6 Suppl 1:S156-162. Pubmed:16534746.
- Revilla, P; Boyat, A; Alvarez, A; Gouesnard, B; Soengas, P; Ordas, A; Malvar, RA, 2006. Heterotic patterns among French and Spanish maize populations. *Maydica* 51:525-535. Pubmed:20070123.
- Rezende, GL; Logullo, C; Meyer, L; Machado, LB; Oliveira-Carvalho, AL; Zingali, RB; Cifuentes, D; Galina, A, 2006. Partial purification of tightly bound mitochondrial hexokinase from maize (*Zea mays* L.) root membranes. *Braz. J. Med. Biol. Res.* 39:1159-1169. Pubmed:16981044.
- Ribaldo, CM; Rondanini, DP; Trinchero, GD; Cura, JA, 2006. Effect of *Herbaspirillum seropedicae* inoculation on maize nitrogen metabolism. *Maydica* 51:481-485. Pubmed:20070123.
- Ribaut, JM; Ragot, M, 2007. Marker-assisted selection to improve drought adaptation in maize: the backcross approach, perspectives, limitations, and alternatives. *J. Exp. Bot.* 58:351-360. Pubmed:17158111.
- Rice, EB; Smith, ME; Mitchell, SE; Kresovich, S, 2006. Conservation and change: A comparison of in situ and ex situ conservation of jala maize germplasm. *Crop Sci.* 46:428-436. Pubmed:20060524.
- Riddle, NC; Kato, A; Birchler, JA, 2006. Genetic variation for the response to ploidy change in *Zea mays* L. *Theor. Appl. Genet.* 114:101-111. Pubmed:17053922.
- Robertson, LA; Kleinschmidt, CE; White, DG; Payne, GA; Maragos, CM; Holland, JB, 2006. Heritabilities and correlations of *Fusarium* ear rot resistance and fumonisin contamination resistance in two maize populations. *Crop Sci.* 46:353-361. Pubmed:20060524.
- Robertson, LA; Kleinschmidt, CE; White, DG; Payne, GA; Maragos, CM; Holland, JB, 2006. Erratum for "Heritabilities and correlations of *Fusarium* ear rot resistance and fumonisin contamination resistance in two maize populations" (vol 46, pg 353, 2006). *Crop Sci.* 46:1420. Pubmed:20060918.
- Robertson-Hoyt, LA; Jines, MP; Balint-Kurti, PJ; Kleinschmidt, CE; White, DG; Payne, GA; Maragos, CM; Molnar, TL; Holland, JB, 2006. QTL mapping for *Fusarium* ear rot and fumonisin contamination resistance in two maize populations. *Crop Sci.* 46:1734-1743. Pubmed:20061017.
- Roda, A; Mirasoli, M; Guardigli, M; Michelini, E; Simoni, P; Magliulo, M, 2006. Development and validation of a sensitive and fast chemiluminescent enzyme immunoassay for the detection of genetically modified maize. *Anal. Bioanal. Chem.* 384:1269-1275. Pubmed:16491341.
- Rodriguez, AA; Ramiro Lascano, H; Bustos, D; Taleisnik, E, 2007. Salinity-induced decrease in NADPH oxidase activity in the maize leaf blade elongation zone. *J. Plant Physiol.* 164:223-230. Pubmed:17074408.
- Rodriguez-F, JG; Sanchez-G, JJ; Baltazar-M, B; De-la-Cruz-L, L; Santacruz-Ruvalcaba, F; Ron-P, J; Schoper, JB, 2006. Characterization of floral morphology and synchrony among *Zea* species in Mexico. *Maydica* 51:383-398. Pubmed:20061003.
- Rodriguez-Nogales, JM; Garcia, MC; Marina, ML, 2006. Analysis of European and North American maize inbred and hybrid lines by monolithic and perfusion reversed-phase high-performance chromatography and multivariate analysis. *J. Agric. Food Chem.* 54:8702-8709. Pubmed:17090110.
- Romanowska, E; Drozak, A, 2006. Comparative analysis of biochemical properties of mesophyll and bundle sheath chloroplasts from various subtypes of C(4) plants grown at moderate irradiance. *Acta Biochim. Pol.* 53:709-719. Pubmed:17106510.
- Romanowska, E; Drozak, A; Pokorska, B; Shiell, BJ; Michalski, WP, 2006. Organization and activity of photosystems in the mesophyll and bundle sheath chloroplasts of maize. *J. Plant Physiol.* 163:607-618. Pubmed:16545994.
- Ross, AJ; Hallauer, AR; Lee, M, 2006. Genetic analysis of traits correlated with maize ear length. *Maydica* 51:301-313. Pubmed:20061003.
- Rossini, L; Vecchietti, A; Nicoloso, L; Stein, N; Franzago, S; Salamini, F; Pozzi, C, 2006. Candidate genes for barley mutants involved in plant architecture: an *in silico* approach. *Theor. Appl. Genet.* 112:1073-1085. Pubmed:16501940.
- Roy-Barman, S; Sautter, C; Chattoo, BB, 2006. Expression of the lipid transfer protein Ace-AMP1 in transgenic wheat enhances antifungal activity and defense responses. *Transgenic Res.* 15:435-446. Pubmed:16906444.

- Royo, J; Gomez, E; Balandin, M; Muniz, LM; Hueros, G, 2006. *ZmLrk-1*, a receptor-like kinase induced by fungal infection in germinating seeds. *Planta* 223:1303-1314. Pubmed:16362327.
- Rudenko, GN; Ono, A; Walbot, V, 2006. An early excision variant of the *MUDR/MU* transposon family is not associated with a local duplication of the *bz1 :: Mu1* allele. *Maydica* 51:227-231. Pubmed:20061003.
- Russell, WK, 2006. Registration of 14 populations of maize developed from a long-term replicated selection study. *Crop Sci.* 46:1824-1825. Pubmed:20061017.
- Russell, WK, 2006. Registration of KLS\_30 and KSS\_30 populations of maize. *Crop Sci.* 46:1405-1406. Pubmed:20060918.
- Russell, WK, 2006. Registration of N551 and N552 parental inbred lines of maize. *Crop Sci.* 46:1014-1015. Pubmed:20060718.
- Sabelli, PA; Larkins, BA, 2006. Grasses like mammals? Redundancy and compensatory regulation within the retinoblastoma protein family. *Cell Cycle* 5:352-355. Pubmed:16479170.
- Sabry, A; Jeffers, D; Vasal, SK; Frederiksen, R; Magill, C, 2006. A region of maize chromosome 2 affects response to downy mildew pathogens. *Theor. Appl. Genet.* 113:321-330. Pubmed:16791698.
- Saccaro, NL, Jr.; Van Sluys, MA; de Mello Varani, A; Rossi, M, 2007. *MudrA*-like sequences from rice and sugarcane cluster as two bona fide transposon clades and two domesticated transposases. *Gene* 392:117-125. Pubmed:17289300.
- Saitoh, T; Ikegami, T; Nakayama, M; Teshima, K; Akutsu, H; Hase, T, 2006. NMR study of the electron transfer complex of plant ferredoxin and sulfite reductase: mapping the interaction sites of ferredoxin. *J. Biol. Chem.* 281:10482-10488. Pubmed:16469743.
- Sala, RG; Andrade, FH; Camadro, EL; Cerono, JC, 2006. Quantitative trait loci for grain moisture at harvest and field grain drying rate in maize (*Zea mays*, L.). *Theor. Appl. Genet.* 112:462-471. Pubmed:16311725.
- Saleh, A; Lumbrieras, V; Lopez, C; Dominguez-Puigjaner, E; Kizis, D; Pages, M, 2006. Maize DBF1-interactor protein 1 containing an R3H domain is a potential regulator of DBF1 activity in stress responses. *Plant J.* 46:747-757. Pubmed:16709191.
- Salhuana, W; Pollak, L, 2006. Latin American Maize Project (LAMP) and Germplasm Enhancement of Maize (GEM) project: Generating useful breeding germplasm. *Maydica* 51:339-355. Pubmed:20061003.
- Sanchez-G, JJ; Goodman, MM; Bird, RM; Stuber, CW, 2006. Isozyme and morphological variation in maize of five Andean countries. *Maydica* 51:25-42. Pubmed:20061017.
- Sanguin, H; Remenant, B; Dechesne, A; Thioulouse, J; Vogel, TM; Nesme, X; Moenne-Loccoz, Y; Grundmann, GL, 2006. Potential of a 16S rRNA-based taxonomic microarray for analyzing the rhizosphere effects of maize on *Agrobacterium* spp. and bacterial communities. *Appl. Environ. Microbiol.* 72:4302-4312. Pubmed:16751545.
- Sanguineti, MC; Duvick, DN; Smith, S; Landi, P; Tuberrosa, R, 2006. Effects of long-term selection on seedling traits and ABA accumulation in commercial maize hybrids. *Maydica* 51:329-338. Pubmed:20061003.
- Santiago, R; Butron, A; Arnason, JT; Reid, LM; Souto, XC; Malvar, RA, 2006. Putative role of pith cell wall phenylpropanoids in *Sesamia nonagrioides* (Lepidoptera: Noctuidae) resistance. *J. Agric. Food Chem.* 54:2274-2279. Pubmed:16536607.
- Santiago, R; Butron, A; Reid, LM; Arnason, JT; Sandoya, G; Souto, XC; Malvar, RA, 2006. Diferulate content of maize sheaths is associated with resistance to the Mediterranean corn borer *Sesamia nonagrioides* (Lepidoptera: Noctuidae). *J. Agric. Food Chem.* 54:9140-9144. Pubmed:17117802.
- Satoh-Nagasawa, N; Nagasawa, N; Malcomber, S; Sakai, H; Jackson, D, 2006. A trehalose metabolic enzyme controls inflorescence architecture in maize. *Nature* 441:227-230. Pubmed:16688177.
- Sauer, M; Jakob, A; Nordheim, A; Hochholdinger, F, 2006. Proteomic analysis of shoot-borne root initiation in maize (*Zea mays* L.). *Proteomics* 6:2530-2541. Pubmed:16521151.
- Sawers, RJ; Farmer, PR; Moffett, P; Brutnell, TP, 2006. In planta transient expression as a system for genetic and biochemical analyses of chlorophyll biosynthesis. *Plant Methods* 2:15. Pubmed:16953878.
- Sawers, RJ; Viney, J; Farmer, PR; Bussey, RR; Olsefski, G; Anufrikova, K; Hunter, CN; Brutnell, TP, 2006. The maize *Oil yellow1* (*Oy1*) gene encodes the I subunit of magnesium chelatase. *Plant Mol. Biol.* 60:95-106. Pubmed:16463102.
- Scaroni, L; Quagliarini, E; Del Buono, D, 2006. Induction of wheat and maize glutathione S-transferase by some herbicide safeners and their effect on enzyme activity against butachlor and terbutylazine. *Pest Manag. Sci.* 62:927-932. Pubmed:16835885.
- Schaeffer, M; Byrne, P; Coe, EH, Jr., 2006. Consensus quantitative trait maps in maize: A database strategy. *Maydica* 51:357-367. Pubmed:20061003.
- Scharff, LB; Koop, HU, 2006. Linear molecules of tobacco ptDNA end at known replication origins and additional loci. *Plant Mol. Biol.* 62:611-621. Pubmed:16897466.
- Schibler, L; Roig, A; Mahe, MF; Laurent, P; Hayes, H; Rodolphe, F; Cribiu, EP, 2006. High-resolution comparative mapping among man, cattle and mouse suggests a role for repeat sequences in mammalian genome evolution. *BMC Genomics* 7:194. Pubmed:16882342.
- Schmitz-Linneweber, C; Williams-Carrier, RE; Williams-Voelker, PM; Kroeger, TS; Vichas, A; Barkan, A, 2006. A pentatricopeptide repeat protein facilitates the trans-splicing of the maize chloroplast *rps12* pre-mRNA. *Plant Cell* 18:2650-2663. Pubmed:17041147.
- Schnee, C; Kollner, TG; Held, M; Turlings, TC; Gershenzon, J; Degenhardt, J, 2006. The products of a single maize sesquiterpene synthase form a volatile defense signal that attracts natural enemies of maize herbivores. *Proc. Natl. Acad. Sci. USA* 103:1129-1134. Pubmed:16418295.
- Schopfer, P; Liszkay, A, 2006. Plasma membrane-generated reactive oxygen intermediates and their role in cell growth of plants. *Biofactors* 28:73-81. Pubmed:17379938.

- Schrag, TA; Melchinger, AE; Sorensen, AP; Frisch, M, 2006. Prediction of single-cross hybrid performance for grain yield and grain dry matter content in maize using AFLP markers associated with QTL. *Theor. Appl. Genet.* 113:1037-1047. Pubmed:16896712.
- Scott, MP; Edwards, JW; Bell, CP; Schussler, JR; Smith, JS, 2006. Grain composition and amino acid content in maize cultivars representing 80 years of commercial maize varieties. *Maydica* 51:417-423. Pubmed:20061003.
- Scott, RJ; Spielman, M, 2006. Deeper into the maize: new insights into genomic imprinting in plants. *Bioessays* 28:1167-1171. Pubmed:17120227.
- Sekhon, RS; Peterson, T; Chopra, S, 2007. Epigenetic modifications of distinct sequences of the *p1* regulatory gene specify tissue-specific expression patterns in maize. *Genetics* 175:1059-1070. Pubmed:17179091.
- Senna, R; Simonin, V; Silva-Neto, MA; Fialho, E, 2006. Induction of acid phosphatase activity during germination of maize (*Zea mays*) seeds. *Plant Physiol. Biochem.* 44:467-473. Pubmed:17023171.
- Shahroona, B; Arshad, M; Zahir, ZA, 2006. Effect of plant growth promoting rhizobacteria containing ACC-deaminase on maize (*Zea mays* L.) growth under axenic conditions and on nodulation in mung bean (*Vigna radiata* L.). *Lett. Appl. Microbiol.* 42:155-159. Pubmed:16441381.
- Shaver, JM; Oldenburg, DJ; Bendich, AJ, 2006. Changes in chloroplast DNA during development in tobacco, *Medicago truncatula*, pea, and maize. *Planta* 224:72-82. Pubmed:16362324.
- She, CW; Liu, JY; Song, YC, 2006. CPD staining: an effective technique for detection of NORs and other GC-rich chromosomal regions in plants. *Biotech. Histochem.* 81:13-21. Pubmed:16760123.
- She, CW; Song, YC, 2006. [Advances in research of the structure and function of plant centromeres]. *Yi Chuan* 28:1597-1606. Pubmed:17138549.
- Sheehan, MJ; Kennedy, LM; Costich, DE; Brutnell, TP, 2007. Subfunctionalization of *PhyB1* and *PhyB2* in the control of seedling and mature plant traits in maize. *Plant J.* 49:338-353. Pubmed:17181778.
- Shenton, M; Fontaine, V; Hartwell, J; Marsh, JT; Jenkins, GI; Nimmo, HG, 2006. Distinct patterns of control and expression amongst members of the *PEP* carboxylase kinase gene family in C4 plants. *Plant J.* 48:45-53. Pubmed:16925599.
- Shepherd, DN; Martin, DP; Varsani, A; Thomson, JA; Rybicki, EP; Klump, HH, 2006. Restoration of native folding of single-stranded DNA sequences through reverse mutations: an indication of a new epigenetic mechanism. *Arch. Biochem. Biophys.* 453:108-122. Pubmed:16427599.
- Sheridan, WF; Auger, DL, 2006. Construction and uses of new compound B-A-A maize chromosome translocations. *Genetics* 174:1755-1765. Pubmed:17057247.
- Shi, C; Koch, G; Ouzunova, M; Wenzel, G; Zein, I; Lubberstedt, T, 2006. Comparison of maize brown-midrib isogenic lines by cellular UV-microspectrophotometry and comparative transcript profiling. *Plant Mol. Biol.* 62:697-714. Pubmed:17016741.
- Shi, J; Dawe, RK, 2006. Partitioning of the maize epigenome by the number of methyl groups on histone H3 lysines 9 and 27. *Genetics* 173:1571-1583. Pubmed:16624902.
- Shim, YS; Kasha, KJ; Simion, E; Letarte, J, 2006. The relationship between induction of embryogenesis and chromosome doubling in microspore cultures. *Protoplasma* 228:79-86. Pubmed:16937058.
- Shimomura, S, 2006. Identification of a glycosylphosphatidylinositol-anchored plasma membrane protein interacting with the C-terminus of auxin-binding protein 1: a photoaffinity crosslinking study. *Plant Mol. Biol.* 60:663-677. Pubmed:16649105.
- Shin, JH; Kwon, SJ; Lee, JK; Min, HK; Kim, NS, 2006. Genetic diversity of maize kernel starch-synthesis genes with SNPs. *Genome* 49:1287-1296. Pubmed:17213911.
- Shin, YM; Park, HJ; Yim, SD; Baek, NI; Lee, CH; An, G; Woo, YM, 2006. Transgenic rice lines expressing maize *C1* and *R-S* regulatory genes produce various flavonoids in the endosperm. *Plant Biotechnol. J.* 4:303-315. Pubmed:17147636.
- Shirgurkar, MV; Naik, VB; von Arnold, S; Nadgauda, RS; Clapham, D, 2006. An efficient protocol for genetic transformation and shoot regeneration of turmeric (*Curcuma longa* L.) via particle bombardment. *Plant Cell Rep.* 25:112-116. Pubmed:16397786.
- Shishova, M; Yemelyanov, V; Rudashevskaya, E; Lindberg, S, 2006. A shift in sensitivity to auxin within development of maize seedlings. *J. Plant Physiol. (in press)*. Pubmed:17074416.
- Shrawat, AK; Lorz, H, 2006. *Agrobacterium*-mediated transformation of cereals: a promising approach crossing barriers. *Plant Biotechnol. J.* 4:575-603. Pubmed:17309731.
- Sidorov, V; Gilbertson, L; Addae, P; Duncan, D, 2006. *Agrobacterium*-mediated transformation of seedling-derived maize callus. *Plant Cell Rep.* 25:320-328. Pubmed:16252091.
- Sineshchekov, VA, 2006. Extreme dehydration of plant tissues irreversibly converts the major and variable phyA' into the minor and conserved phyA". *J. Photochem. Photobiol. B* 85:85-91. Pubmed:16829116.
- Singh, J; Zhang, S; Chen, C; Cooper, L; Bregitzer, P; Sturbaum, A; Hayes, PM; Lemaux, PG, 2006. High-frequency *Ds* remobilization over multiple generations in barley facilitates gene tagging in large genome cereals. *Plant Mol. Biol.* 62:937-950. Pubmed:17004014.
- Sivakumar, S; Franco, OL; Thayumanavan, B; Murad, AM; Manickam, A; Mohan, M; Mridula, M, 2006. Cloning and structural analysis of an Indian little millet (*Panicum sumatrense*) zein-like storage protein: implications for molecular assembly. *Biochemistry (Mosc.)* 71:1183-1191. Pubmed:17140379.
- Sivamani, E; Qu, R, 2006. Expression enhancement of a rice polyubiquitin gene promoter. *Plant Mol. Biol.* 60:225-239. Pubmed:16429261.

- Sluyter, A; Dominguez, G, 2006. Early maize (*Zea mays* L.) cultivation in Mexico: dating sedimentary pollen records and its implications. Proc. Natl. Acad. Sci. USA 103:1147-1151. Pubmed:16418287.
- Smidansky, ED; Meyer, FD; Blakeslee, B; Weglarz, TE; Greene, TW; Giroux, MJ, 2006. Expression of a modified ADP-glucose pyrophosphorylase large subunit in wheat seeds stimulates photosynthesis and carbon metabolism. Planta (in press). Pubmed:17021802.
- Smith, JSC; Desbons, P; Gogerty, J; Niebur, WS, 2006. Changes in parentage and genetic diversity of widely used maize hybrids grown in the northern United States and France from 1930 to the present. Maydica 51:57-77. Pubmed:20061017.
- Smith, S; Loffler, C; Cooper, M, 2006. Genetic diversity among maize hybrids widely grown in contrasting regional environments in the United States during the 1990s. Maydica 51:233-242. Pubmed:20061003.
- Soengas, P; Ordas, B; Malvar, R-A; Revilla, P; Ordas, A, 2006. Combining abilities and heterosis for adaptation in flint maize populations. Crop Sci. 46:2666-2669. Pubmed:20070306.
- Sokolov, VA, 2006. [Imprinting in plants]. Genetika 42:1250-1260. Pubmed:17100092.
- Sparg, SG; Kulkarni, MG; van-Staden, J, 2006. Aerosol smoke and smoke-water stimulation of seedling vigor of a commercial maize cultivar. Crop Sci. 46:1336-1340. Pubmed:20060918.
- Spielbauer, G; Margl, L; Hannah, LC; Romisch, W; Ettenhuber, C; Bacher, A; Gierl, A; Eisenreich, W; Genschel, U, 2006. Robustness of central carbohydrate metabolism in developing maize kernels. Phytochemistry 67:1460-1475. Pubmed:16815503.
- Spitko, T; Sagi, L; Pinter, J; Marton, LC; Barnabas, B, 2006. Haploid regeneration aptitude of maize (*Zea mays* L.) lines of various origin and of their hybrids. Maydica 51:537-542. Pubmed:20070123.
- Stass, A; Wang, Y; Eticha, D; Horst, WJ, 2006. Aluminium rhizotoxicity in maize grown in solutions with Al<sup>3+</sup> or Al(OH)<sup>-4</sup> as predominant solution Al species. J. Exp. Bot. 57:4033-4042. Pubmed:17105968.
- Stich, B; Melchinger, AE; Piepho, HP; Heckenberger, M; Maurer, HP; Reif, JC, 2006. A new test for family-based association mapping with inbred lines from plant breeding programs. Theor. Appl. Genet. 113:1121-1130. Pubmed:16896707.
- Stich, B; Yu, J; Melchinger, AE; Piepho, HP; Utz, F; Maurer, HP; Buckler, ES, 2006. Power to detect higher-order epistatic interactions in a metabolic pathway using a new mapping strategy. Genetics (in press). Pubmed:17194777.
- Sticklen, M, 2006. Plant genetic engineering to improve biomass characteristics for biofuels. Curr. Opin. Biotechnol. 17:315-319. Pubmed:16701991.
- Stige, LC; Stave, J; Chan, KS; Ciannelli, L; Pettorelli, N; Glantz, M; Herren, HR; Stenseth, NC, 2006. The effect of climate variation on agro-pastoral production in Africa. Proc. Natl. Acad. Sci. USA 103:3049-3053. Pubmed:16492727.
- Storer, NP; Babcock, JM; Edwards, JM, 2006. Field measures of western corn rootworm (Coleoptera: Chrysomelidae) mortality caused by Cry34/35Ab1 proteins expressed in maize event 59122 and implications for trait durability. J. Econ. Entomol. 99:1381-1387. Pubmed:16937696.
- Straska, J; Sebela, M; Tarkowski, P; Rehulka, P; Chmelik, J; Popa, I; Pec, P, 2007. Inhibition of plant amine oxidases by a novel series of diamine derivatives. Biochimie 89:135-144. Pubmed:16989933.
- Stupar, RM; Springer, NM, 2006. Cis-transcriptional variation in maize inbred lines B73 and Mo17 leads to additive expression patterns in the F1 hybrid. Genetics 173:2199-2210. Pubmed:16702414.
- Su, W; Lin, C; Wu, J; Li, K; He, G; Qian, X; Wei, C; Yang, J, 2006. Molecular cloning and expression of a cDNA encoding Lon protease from rice (*Oryza sativa*). Biotechnol. Lett. 28:923-927. Pubmed:16786279.
- Subbaiah, CC; Palaniappan, A; Duncan, K; Rhoads, DM; Huber, SC; Sachs, MM, 2006. Mitochondrial localization and putative signaling function of sucrose synthase in maize. J. Biol. Chem. 281:15625-15635. Pubmed:16606624.
- Subedi, KD; Ma, BL; Smith, DL, 2006. Response of a leafy and non-leafy maize hybrid to population densities and fertilizer nitrogen levels. Crop Sci. 46:1860-1869. Pubmed:20061219.
- Subramanian, S; Bonen, L, 2006. Rapid evolution in sequence and length of the nuclear-located gene for mitochondrial L2 ribosomal protein in cereals. Genome 49:275-281. Pubmed:16604111.
- Sudan, C; Prakash, S; Bhomkar, P; Jain, S; Bhalla-Sarin, N, 2006. Ubiquitous presence of beta-glucuronidase (GUS) in plants and its regulation in some model plants. Planta 224:853-864. Pubmed:16652220.
- Suen, DF; Huang, AH, 2007. Maize pollen coat xylanase facilitates pollen tube penetration into silk during sexual reproduction. J. Biol. Chem. 282:625-636. Pubmed:17062571.
- Sun, B; Jing, Y; Chen, K; Song, L; Chen, F; Zhang, L, 2007. Protective effect of nitric oxide on iron deficiency-induced oxidative stress in maize (*Zea mays*). J. Plant Physiol. 164:536-543. Pubmed:16690167.
- Suzuki, M; Mark Settles, A; Tsueung, CW; Li, QB; Latshaw, S; Wu, S; Porch, TG; Schmelz, EA; James, MG; McCarty, DR, 2006. The maize *viviparous15* locus encodes the molybdopterin synthase small subunit. Plant J. 45:264-274. Pubmed:16367969.
- Swanson-Wagner, RA; Jia, Y; DeCook, R; Borsuk, LA; Nettleton, D; Schnable, PS, 2006. All possible modes of gene action are observed in a global comparison of gene expression in a maize F1 hybrid and its inbred parents. Proc. Natl. Acad. Sci. USA 103:6805-6810. Pubmed:16641103.
- Sweeney, MT; Thomson, MJ; Pfeil, BE; McCouch, S, 2006. Caught red-handed: *Rc* encodes a basic helix-loop-helix protein conditioning red pericarp in rice. Plant Cell 18:283-294. Pubmed:16399804.
- Taba, S; Diaz, J; Aragon-C, F; Rincon-Sanchez, F; Hernandez, JM; Krakowsky, M, 2006. Evaluation of Zapalote Chico accessions for conservation and enhancement. Maydica 51:209-218. Pubmed:20061003.

- Tan, S; Evans, R; Singh, B, 2006. Herbicidal inhibitors of amino acid biosynthesis and herbicide-tolerant crops. *Amino Acids* 30:195-204. Pubmed:16547651.
- Tang, JH; Fu, ZY; Hu, YM; Li, JS; Sun, LL; Ji, HQ, 2006. Genetic analyses and mapping of a new thermo-sensitive genic male sterile gene in maize. *Theor. Appl. Genet.* 113:11-15. Pubmed:16783588.
- Tang, W; Coughlan, S; Crane, E; Beatty, M; Duvick, J, 2006. The application of laser microdissection to in planta gene expression profiling of the maize anthracnose stalk rot fungus *Colletotrichum graminicola*. *Mol. Plant Microbe Interact.* 19:1240-1250. Pubmed:17073306.
- Tao, F; Zhu, SW; Fan, J; Cheng, BJ, 2006. Cloning and sequence analysis of maize *FAD2* gene. *Zhi Wu Sheng Li Yu Fen Zi Sheng Wu Xue Xue Bao* 32:649-656. Pubmed:17167201.
- Tao, TY; Ouellet, T; Dadej, K; Miller, SS; Johnson, DA; Singh, J, 2006. Characterization of a novel glycine-rich protein from the cell wall of maize silk tissues. *Plant Cell Rep.* 25:848-858. Pubmed:16528565.
- Tarter, JA; Holland, JB, 2006. Gains from selection during the development of semixotic inbred lines from Latin American maize accessions. *Maydica* 51:15-23. Pubmed:20061017.
- Tatum, TC; Rayburn, AL, 2006. PRINS-labeled knobs are not associated with increased chromosomal stickiness in the maize *st1* mutant. *J. Hered.* 97:417-422. Pubmed:16837564.
- Tavladoraki, P; Rossi, MN; Saccuti, G; Perez-Amador, MA; Polticelli, F; Angelini, R; Federico, R, 2006. Heterologous expression and biochemical characterization of a polyamine oxidase from *Arabidopsis* involved in polyamine back conversion. *Plant Physiol.* 141:1519-1532. Pubmed:16778015.
- Teardo, E; de Laureto, PP; Bergantino, E; Dalla Vecchia, F; Rigoni, F; Szabo, I; Giacometti, GM, 2006. Evidences for interaction of PsbS with photosynthetic complexes in maize thylakoids. *Biochim. Biophys. Acta* (in press). Pubmed:17250801.
- Teshima, KM; Coop, G; Przeworski, M, 2006. How reliable are empirical genomic scans for selective sweeps? *Genome Res.* 16:702-712. Pubmed:16687733.
- Thompson, D-L; Goodman, M-M, 2006. Increasing kernel density for two inbred lines of maize. *Crop Sci.* 46:2179-2182. Pubmed:20061219.
- Tiffin, P, 2006. Population genetic evidence for rapid changes in intra-specific diversity and allelic cycling of a specialist defense gene in *Zeae*. *Genetics* (in press). Pubmed:16582423.
- Tobias, DJ; Manoharan, M; Pritsch, C; Dahleen, LS, 2007. Co-bombardment, integration and expression of rice chitinase and thaumatin-like protein genes in barley (*Hordeum vulgare* cv. Conlon). *Plant Cell Rep.* 26:631-639. Pubmed:17103003.
- Tollenaar, M; Lee, EA, 2006. Dissection of physiological processes underlying grain yield in maize by examining genetic improvement and heterosis. *Maydica* 51:399-408. Pubmed:20061003.
- Tomilov, A; Tomilova, N; Yoder, JI, 2007. *Agrobacterium tumefaciens* and *Agrobacterium* rhizogenes transformed roots of the parasitic plant *Triphysaria versicolor* retain parasitic competence. *Planta* 225:1059-1071. Pubmed:17053892.
- Ton, J; D'Alessandro, M; Jourdie, V; Jakab, G; Karlen, D; Held, M; Mauch-Mani, B; Turlings, TC, 2007. Priming by airborne signals boosts direct and indirect resistance in maize. *Plant J.* 49:16-26. Pubmed:17144894.
- Toyota, A; Akiyama, H; Sugimura, M; Watanabe, T; Sakata, K; Shiramasa, Y; Kitta, K; Hino, A; Esaka, M; Maitani, T, 2006. Rapid quantification methods for genetically modified maize contents using genomic DNAs pretreated by sonication and restriction endonuclease digestion for a capillary-type real-time PCR system with a plasmid reference standard. *Biosci. Biotechnol. Biochem.* 70:2965-2973. Pubmed:17151472.
- Tracy, WF; Whitt, SR; Buckler, ES, 2006. Recurrent mutation and genome evolution: Example of *Sugary 1* and the origin of sweet maize. *Crop Sci.* 46:S49-S54. Pubmed:20070306.
- Tsugane, K; Maekawa, M; Takagi, K; Takahara, H; Qian, Q; Eun, CH; Iida, S, 2006. An active DNA transposon *nDart* causing leaf variegation and mutable dwarfism and its related elements in rice. *Plant J.* 45:46-57. Pubmed:16367953.
- Uchibayashi, M, 2006. [Presence of pre-Columbian maize in the Old World—an overview]. *Yakugaku Zasshi* 126:423-427. Pubmed:16755129.
- Uchibayashi, M, 2006. [Maize in pre-columbian China found in Bencao Pinhui Jingyao]. *Yakugaku Zasshi* 126:27-36. Pubmed:16394647.
- Unno, H; Uchida, T; Sugawara, H; Kurisu, G; Sugiyama, T; Yamaya, T; Sakakibara, H; Hase, T; Kusunoki, M, 2006. Atomic structure of plant glutamine synthetase: a key enzyme for plant productivity. *J. Biol. Chem.* 281:29287-29296. Pubmed:16829528.
- Upadyayula, N; da Silva, HS; Bohn, MO; Rocheford, TR, 2006. Genetic and QTL analysis of maize tassel and ear inflorescence architecture. *Theor. Appl. Genet.* 112:592-606. Pubmed:16395569.
- Upadyayula, N; Wassom, J; Bohn, MO; Rocheford, TR, 2006. Quantitative trait loci analysis of phenotypic traits and principal components of maize tassel inflorescence architecture. *Theor. Appl. Genet.* 113:1395-1407. Pubmed:17061102.
- Uzarowska, A; Keller, B; Piepho, HP; Schwarz, G; Ingvarsson, C; Wenzel, G; Lubberstedt, T, 2007. Comparative expression profiling in meristems of inbred-hybrid triplets of maize based on morphological investigations of heterosis for plant height. *Plant Mol. Biol.* 63:21-34. Pubmed:17006594.
- van Etten, J, 2006. Changes in farmers' knowledge of maize diversity in highland Guatemala, 1927/37-2004. *J. Ethnobiol. Ethnomedicine* 2:12. Pubmed:16507113.
- van Leeuwen, H; Monfort, A; Puigdomenech, P, 2007. Mutator-like elements identified in melon, *Arabidopsis* and rice contain ULP1 protease domains. *Mol. Genet. Genomics* 277:357-364. Pubmed:17136348.

- Vargas, M; van Eeuwijk, FA; Crossa, J; Ribaut, JM, 2006. Mapping QTLs and QTL x environment interaction for CIMMYT maize drought stress program using factorial regression and partial least squares methods. *Theor. Appl. Genet.* 112:1009-1023. Pubmed:16538513.
- Vasic, NJ; Ivanovic, MR; Brkic, IJ; Bekavac, GF; Zdunec, ZI; Jambrovic, AS, 2006. Evaluation of maize hybrids containing different proportion of NC298 tropical germplasm line in their male parents. *Maydica* 51:79-88. Pubmed:20061017.
- Vasil, IK; Vasil, V, 2006. Transformation of wheat via particle bombardment. *Methods Mol. Biol.* 318:273-283. Pubmed:16673923.
- Vij, S; Gupta, V; Kumar, D; Vydiathan, R; Raghuvanshi, S; Khurana, P; Khurana, JP; Tyagi, AK, 2006. Decoding the rice genome. *Bioessays* 28:421-432. Pubmed:16547947.
- Vinagre, F; Vargas, C; Schwarcz, K; Cavalcante, J; Nogueira, EM; Baldani, JL; Ferreira, PC; Hemerly, AS, 2006. SHR5: a novel plant receptor kinase involved in plant-N2-fixing endophytic bacteria association. *J. Exp. Bot.* 57:559-569. Pubmed:16397001.
- Vitte, C; Bennetzen, JL, 2006. Analysis of retrotransposon structural diversity uncovers properties and propensities in angiosperm genome evolution. *Proc. Natl. Acad. Sci. USA* 103:17638-17643. Pubmed:17101966.
- Voisin, AS; Reidy, B; Parent, B; Rolland, G; Redondo, E; Gerentes, D; Tardieu, F; Muller, B, 2006. Are ABA, ethylene or their interaction involved in the response of leaf growth to soil water deficit? An analysis using naturally occurring variation or genetic transformation of ABA production in maize. *Plant Cell Environ.* 29:1829-1840. Pubmed:16913872.
- Walker, NS; Stiffler, N; Barkan, A, 2007. POGs/PlantRBP: a resource for comparative genomics in plants. *Nucl. Acids Res.* 35:D852-856. Pubmed:17142226.
- Walter, MH; Floss, DS; Hans, J; Fester, T; Strack, D, 2007. Apocarotenoid biosynthesis in arbuscular mycorrhizal roots: contributions from methylerythritol phosphate pathway isogenes and tools for its manipulation. *Phytochemistry* 68:130-138. Pubmed:17084869.
- Walton, JD, 2006. HC-toxin. *Phytochemistry* 67:1406-1413. Pubmed:16839576.
- Wang, BB; Brendel, V, 2006. Molecular characterization and phylogeny of U2AF35 homologs in plants. *Plant Physiol.* 140:624-636. Pubmed:16407443.
- Wang, CJ; Harper, L; Cande, WZ, 2006. High-resolution single-copy gene fluorescence in situ hybridization and its use in the construction of a cytogenetic map of maize chromosome 9. *Plant Cell* 18:529-544. Pubmed:16461583.
- Wang, JW; Yang, FP; Chen, XQ; Liang, RQ; Zhang, LQ; Geng, DM; Zhang, XD; Song, YZ; Zhang, GS, 2006. Induced expression of DREB transcriptional factor and study on its physiological effects of drought tolerance in transgenic wheat. *Yi Chuan Xue Bao* 33:468-476. Pubmed:16722342.
- Wang, Q; Dooner, HK, 2006. Remarkable variation in maize genome structure inferred from haplotype diversity at the bz locus. *Proc. Natl. Acad. Sci. USA* 103:17644-17649. Pubmed:17101975.
- Wang, W; Milanesi, C; Falieri, C; Cresti, M, 2006. Localization of group-1 allergen Zea m 1 in the coat and wall of maize pollen. *Acta Histochem.* 108:395-400. Pubmed:16963110.
- Wang, W; Zheng, H; Fan, C; Li, J; Shi, J; Cai, Z; Zhang, G; Liu, D; Zhang, J; Vang, S; Lu, Z; Wong, GK; Long, M; Wang, J, 2006. High rate of chimeric gene origination by retroposition in plant genomes. *Plant Cell* 18:1791-1802. Pubmed:16829590.
- Watanabe, T; Kasama, K; Kikuchi, H; Suzuki, T; Tokishita, S; Sakata, K; Matsuki, A; Hino, A; Akiyama, H; Maitani, T, 2006. [Laboratory-performance study of quantitative PCR methods to analyze an approved genetically modified maize (Mon810 Line)]. *Shokuhin Eiseigaku Zasshi* 47:15-27. Pubmed:16619852.
- Waters, BM; Chu, HH; Didonato, RJ; Roberts, LA; Eisley, RB; Lahner, B; Salt, DE; Walker, EL, 2006. Mutations in *Arabidopsis yellow stripe-like1* and *yellow stripe-like3* reveal their roles in metal ion homeostasis and loading of metal ions in seeds. *Plant Physiol.* 141:1446-1458. Pubmed:16815956.
- Weekes, R; Allnutt, T; Boffey, C; Morgan, S; Bilton, M; Daniels, R; Henry, C, 2007. A study of crop-to-crop gene flow using farm scale sites of fodder maize (*Zea mays* L.) in the UK. *Transgenic Res.* 16:203-211. Pubmed:17115253.
- Weinstock, BA; Janni, J; Hagen, L; Wright, S, 2006. Prediction of oil and oleic acid concentrations in individual corn (*Zea mays* L.) kernels using near-infrared reflectance hyperspectral imaging and multivariate analysis. *Appl. Spectrosc.* 60:9-16. Pubmed:16454902.
- Welcker, C; Boussuge, B; Bencivenni, C; Ribaut, JM; Tardieu, F, 2007. Are source and sink strengths genetically linked in maize plants subjected to water deficit? A QTL study of the responses of leaf growth and of anthesis-silking interval to water deficit. *J. Exp. Bot.* 58:339-349. Pubmed:17130185.
- Wendl, MC, 2006. A general coverage theory for shotgun DNA sequencing. *J. Comput. Biol.* 13:1177-1196. Pubmed:16901236.
- Weng, LX; Deng, H; Xu, JL; Li, Q; Wang, LH; Jiang, Z; Zhang, HB; Zhang, LH, 2006. Regeneration of sugarcane elite breeding lines and engineering of stem borer resistance. *Pest Manag. Sci.* 62:178-187. Pubmed:16408322.
- Wenkai, X; Mingliang, X; Jiuren, Z; Fengge, W; Jiansheng, L; Jingrui, D, 2006. Genome-wide isolation of resistance gene analogs in maize (*Zea mays* L.). *Theor. Appl. Genet.* 113:63-72. Pubmed:16607513.
- Wenzel, G, 2006. Molecular plant breeding: achievements in green biotechnology and future perspectives. *Appl. Microbiol. Biotechnol.* 70:642-650. Pubmed:16586104.
- West, DR; Kincer, DR; Allen, FL; Thompson, MA, 2006. Registration of maize parental line T175. *Crop Sci.* 46:2733-2734. Pubmed:20070306.
- Whitehead, FC; Caton, HG; Hallauer, AR; Vasal, S; Cordova, H, 2006. Incorporation of elite subtropical and tropical maize germplasm into elite temperate germplasm. *Maydica* 51:43-56. Pubmed:20061017.
- Wight, CP; Kibite, S; Tinker, NA; Molnar, SJ, 2006. Identification of molecular markers for aluminium tolerance in diploid oat through comparative mapping and QTL analysis. *Theor. Appl. Genet.* 112:222-231. Pubmed:16323000.

- Williams, P; Hardeman, K; Fowler, J; Rivin, C, 2006. Divergence of duplicated genes in maize: evolution of contrasting targeting information for enzymes in the porphyrin pathway. *Plant J.* 45:727-739. Pubmed:16460507.
- Williams, WP; Windham, GL, 2006. Registration of maize germplasm line Mp717. *Crop Sci.* 46:1407-1408. Pubmed:20060918.
- Willmot, DB; Dudley, JW; Rocheford, TR; Bari, A, 2006. Effect of random mating on marker-QTL associations for grain quality traits in the cross of Illinois High Oil X Illinois Low Oil. *Maydica* 51:187-199. Pubmed:20061003.
- Wolf, T; Heidelmann, T; Marten, I, 2006. ABA regulation of K(+) -permeable channels in maize subsidiary cells. *Plant Cell Physiol.* 47:1372-1380. Pubmed:16973684.
- Woll, K; Dressel, A; Sakai, H; Piepho, HP; Hochholdinger, F, 2006. ZmGrp3: identification of a novel marker for root initiation in maize and development of a robust assay to quantify allele-specific contribution to gene expression in hybrids. *Theor. Appl. Genet.* 113:1305-1315. Pubmed:16937154.
- Wolt, JD; Conlan, CA; Majima, K, 2005. An ecological risk assessment of Cry1F maize pollen impact to pale grass blue butterfly. *Environ. Biosafety Res.* 4:243-251. Pubmed:16827552.
- Woodhouse, MR; Freeling, M; Lisch, D, 2006. Initiation, establishment, and maintenance of heritable MuDR transposon silencing in maize are mediated by distinct factors. *PLoS Biol.* 4:e339. Pubmed:16968137.
- Woodhouse, MR; Freeling, M; Lisch, D, 2006. The *mop1* (*mediator of paramutation1*) mutant progressively reactivates one of the two genes encoded by the MuDR transposon in maize. *Genetics* 172:579-592. Pubmed:16219782.
- Wu, AH; Zhang, SQ; Deng, XP; Shan, L; Liu, XF, 2006. [Expression of ZmPIP1 subgroup genes in maize roots under water shortage]. *Zhi Wu Sheng Li Yu Fen Zi Sheng Wu Xue Xue Bao* 32:557-562. Pubmed:17075179.
- Wu, XR; Chen, Z; Shende, A; Dooner, HK; Folk, WR, 2006. Visualizing bz1 missense suppression in *Zea mays*: an assay for monocot tRNA expression and utilization. *Plant Mol. Biol.* 61:795-798. Pubmed:16897493.
- Xiao, H; Zhang, F; Zheng, Y, 2006. The 5' stem-loop and its role in mRNA stability in maize S cytoplasmic male sterility. *Plant J.* 47:864-872. Pubmed:16961731.
- Xie, WQ; Zhao, LQ; Bai, WY; Li, ZP; Zhao, YL; Li, CF, 2006. [The effects of calmodulin on the lipid-binding activity of CaM-binding protein-10 and maize non-specific lipid transfer protein]. *Zhi Wu Sheng Li Yu Fen Zi Sheng Wu Xue Xue Bao* 32:679-684. Pubmed:17167205.
- Xing, Y; Ingvarsson, C; Salomon, R; Lubberstedt, T, 2006. Analysis of sugarcane mosaic virus resistance in maize in an isogenic dihybrid crossing scheme and implications for breeding potyvirus-resistant maize hybrids. *Genome* 49:1274-1282. Pubmed:17213909.
- Xiong, GS; Hu, XM; Jiao, YQ; Yu, YC; Chu, CC; Li, JY; Qian, Q; Wang, YH, 2006. Leafy head2, which encodes a putative RNA-binding protein, regulates shoot development of rice. *Cell Res.* 16:267-276. Pubmed:16541125.
- Xu, J; Miao, H; Wu, H; Huang, W; Tang, R; Qiu, M; Wen, J; Zhu, S; Li, Y, 2006. Screening genetically modified organisms using multiplex PCR coupled with oligonucleotide microarray. *Biosens. Bioelectron.* 22:71-77. Pubmed:16466912.
- Xu, JH; Messing, J, 2006. Maize haplotype with a helitron-amplified cytidine deaminase gene copy. *BMC Genet.* 7:52. Pubmed:17094807.
- Xu, W; Ahmed, S; Moriyama, H; Chollet, R, 2006. The importance of the strictly conserved, C-terminal glycine residue in phosphoenolpyruvate carboxylase for overall catalysis: mutagenesis and truncation of GLY-961 in the sorghum C4 leaf isoform. *J. Biol. Chem.* 281:17238-17245. Pubmed:16624802.
- Xu, WH; Zhang, QR; Denlinger, DL, 2006. A novel member of the NSF family in the corn earworm, *Helicoverpa zea*: molecular cloning, developmental expression, and tissue distribution. *Biochim. Biophys. Acta* 1759:186-190. Pubmed:16757382.
- Xu, Z; Dooner, HK, 2006. The maize aberrant pollen transmission 1 gene is a SABRE/KIP homolog required for pollen tube growth. *Genetics* 172:1251-1261. Pubmed:16299389.
- Yamaguchi, T; Lee, DY; Miyao, A; Hirochika, H; An, G; Hirano, HY, 2006. Functional diversification of the two C-class MADS box genes OSMADS3 and OSMADS58 in *Oryza sativa*. *Plant Cell* 18:15-28. Pubmed:16326928.
- Yamane, K; Yano, K; Kawahara, T, 2006. Pattern and rate of indel evolution inferred from whole chloroplast intergenic regions in sugarcane, maize and rice. *DNA Res.* 13:197-204. Pubmed:17110395.
- Yandeau-Nelson, MD; Nikolau, BJ; Schnable, PS, 2006. Effects of trans-acting genetic modifiers on meiotic recombination across the *a1-sh2* interval of maize. *Genetics* 174:101-112. Pubmed:16816431.
- Yandeau-Nelson, MD; Xia, Y; Li, J; Neuffer, MG; Schnable, PS, 2006. Unequal sister chromatid and homolog recombination at a tandem duplication of the *A1* locus in maize. *Genetics* 173:2211-2226. Pubmed:16751673.
- Yang, H; Kaur, N; Kiriakopolos, S; McCormick, S, 2006. EST generation and analyses towards identifying female gametophyte-specific genes in *Zea mays* L. *Planta* 224:1004-1014. Pubmed:16718485.
- Yang, Y; Guo, A; An, S; Liu, G, 2006. [Mathematical simulation of stomatal regulation involving root-sourced signaling ABA under soil drought condition]. *Ying Yong Sheng Tai Xue Bao* 17:65-70. Pubmed:16689236.
- Yang, ZQ; Liu, QQ; Pan, ZM; Yu, HX; Jiao, XA, 2007. Expression of the fusion glycoprotein of Newcastle disease virus in transgenic rice and its immunogenicity in mice. *Vaccine* 25:591-598. Pubmed:17049688.
- Yennawar, NH; Li, LC; Dudzinski, DM; Tabuchi, A; Cosgrove, DJ, 2006. Crystal structure and activities of EXPB1 (*Zea m 1*), a beta-expansin and group-1 pollen allergen from maize. *Proc. Natl. Acad. Sci. USA* 103:14664-14671. Pubmed:16984999.
- Yoke-Kqueen, C; Radu, S, 2006. Random amplified polymorphic DNA analysis of genetically modified organisms. *J. Biotechnol.* 127:161-166. Pubmed:16860900.

- Yu, C; Li, Z, 2006. Construction of methylation-sensitive partial restriction bacterial artificial chromosome libraries in maize. *Anal Biochem.* 359:141-143. Pubmed:17010300.
- Yu, J; Buckler, ES, 2006. Genetic association mapping and genome organization of maize. *Curr. Opin. Biotechnol.* 17:155-160. Pubmed:16504497.
- Yu, J; Pressoir, G; Briggs, WH; Vroh Bi, I; Yamasaki, M; Doebley, JF; McMullen, MD; Gaut, BS; Nielsen, DM; Holland, JB; Kresovich, S; Buckler, ES, 2006. A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. *Nat. Genet.* 38:203-208. Pubmed:16380716.
- Yu, W; Han, F; Kato, A; Birchler, JA, 2006. Characterization of a maize isochromosome 8S\*8S. *Genome* 49:700-706. Pubmed:16936849.
- Yu, W; Lamb, JC; Han, F; Birchler, JA, 2006. Telomere-mediated chromosomal truncation in maize. *Proc. Natl. Acad. Sci. USA* 103:17331-17336. Pubmed:17085598.
- Yu, W; Lamb, JC; Han, F; Birchler, JA, 2007. Cytological visualization of DNA transposons and their transposition pattern in somatic cells of maize. *Genetics* 175:31-39. Pubmed:17057234.
- Yu, XZ; Gu, JD, 2006. Differences in Michaelis-Menten kinetics for different cultivars of maize during cyanide removal. *Ecotoxicol. Environ. Safety* 67:254-259. Pubmed:17064775.
- Yuan, J; Sayegh, J; Mendez, J; Sward, L; Sanchez, N; Sanchez, S; Waldrop, G; Grover, S, 2006. The regulatory role of residues 226-232 in phosphoenolpyruvate carboxylase from maize. *Photosynth. Res.* 88:73-81. Pubmed:16453061.
- Zandonadi, DB; Canellas, LP; Facanha, AR, 2007. Indolacetic and humic acids induce lateral root development through a concerted plasmalemma and tonoplast H<sup>(+)</sup> pumps activation. *Planta* 225:1583-1595. Pubmed:17180358.
- Zhang, A; Jiang, M; Zhang, J; Tan, M; Hu, X, 2006. Mitogen-activated protein kinase is involved in abscisic acid-induced antioxidant defense and acts downstream of reactive oxygen species production in leaves of maize plants. *Plant Physiol.* 141:475-487. Pubmed:16531486.
- Zhang, B; Pan, X; Anderson, TA, 2006. Identification of 188 conserved maize microRNAs and their targets. *FEBS Lett.* 580:3753-3762. Pubmed:16780841.
- Zhang, F; Peterson, T, 2006. Gene conversion between direct noncoding repeats promotes genetic and phenotypic diversity at a regulatory locus of *Zea mays* (L.). *Genetics* 174:753-762. Pubmed:16816430.
- Zhang, F; Wan, XQ; Pan, GT, 2006. QTL mapping of *Fusarium moniliforme* ear rot resistance in maize. 1. Map construction with microsatellite and AFLP markers. *J. Appl. Genet.* 47:9-15. Pubmed:16424603.
- Zhang, J; Zhang, F; Peterson, T, 2006. Transposition of reversed Ac element ends generates novel chimeric genes in maize. *PLoS Genet.* 2:e164. Pubmed:17029561.
- Zhang, LB; Ge, S, 2007. Multilocus analysis of nucleotide variation and speciation in *Oryza officinalis* and its close relatives. *Mol. Biol. Evol.* 24:769-783. Pubmed:17182895.
- Zhang, Y; Wang, L; Liu, Y; Zhang, Q; Wei, Q; Zhang, W, 2006. Nitric oxide enhances salt tolerance in maize seedlings through increasing activities of proton-pump and Na<sup>+</sup>/H<sup>+</sup> antiport in the tonoplast. *Planta* 224:545-555. Pubmed:16501990.
- Zhang, ZF; Wang, Y; Zheng, YL, 2006. AFLP and PCR-based markers linked to *Rf3*, a fertility restorer gene for S cytoplasmic male sterility in maize. *Mol. Genet. Genomics* 276:162-169. Pubmed:16705419.
- Zhang, ZM; Zhao, MJ; Ding, HP; Rong, TZ; Pan, GT, 2006. QTL mapping analysis of plant height and ear height of maize (*Zea mays* L.). *Genetika* 42:391-396. Pubmed:16649666.
- Zhang, ZX; Zhang, SP; Zheng, YL, 2006. [A strategy based on comparative genomics to align ESTs of maize]. *Yi Chuan* 28:339-344. Pubmed:16551603.
- Zhao, HH; Yan, YL, 2006. [The effects of noise on NIR analysis and related mathematic pretreatments and models]. *Guang Pu Xue Yu Guang Pu Fen Xi* 26:842-845. Pubmed:16883850.
- Zhao, M; Zhang, Z; Zhang, S; Li, W; Jeffers, D-P; Rong, T; Pan, G, 2006. Quantitative trait loci for resistance to banded leaf and sheath blight in maize. *Crop Sci.* 46:1039-1045. Pubmed:20060918.
- Zhao, MJ; Gao, SB; Zhang, ZM; Rong, TZ; Pan, GT, 2006. [Initial identification of quantitative trait loci controlling resistance to banded leaf and sheath blight at elongating and heading date in maize]. *Fen Zi Xi Bao Sheng Wu Xue Bao* 39:139-144. Pubmed:16944585.
- Zhao, T; Palotta, M; Langridge, P; Prasad, M; Graner, A; Schulze-Lefert, P; Koprek, T, 2006. Mapped *Ds*/T-DNA launch pads for functional genomics in barley. *Plant J.* 47:811-826. Pubmed:16889649.
- Zhao, W; Canaran, P; Jurkuta, R; Fulton, T; Glaubitz, J; Buckler, E; Doebley, J; Gaut, B; Goodman, M; Holland, J; Kresovich, S; McMullen, M; Stein, L; Ware, D, 2006. PanZeal: a database and resource for molecular and functional diversity in the maize genome. *Nucl. Acids Res.* 34:D752-757. Pubmed:16381974.
- Zhao, ZY; Ranch, J, 2006. Transformation of maize via *Agrobacterium tumefaciens* using a binary co-integrate vector system. *Methods Mol. Biol.* 318:315-323. Pubmed:16673926.
- Zhu, D; Jiang, MY; Tan, MP, 2006. [The mechanism of ABA-induced apoplastic H<sub>2</sub>O<sub>2</sub> accumulation in maize leaves]. *Zhi Wu Sheng Li Yu Fen Zi Sheng Wu Xue Bao* 32:519-526. Pubmed:17075174.
- Zhu, J; Chen, S; Alvarez, S; Asirvatham, VS; Schachtman, DP; Wu, Y; Sharp, RE, 2006. Cell wall proteome in the maize primary root elongation zone. I. Extraction and identification of water-soluble and lightly ionically bound proteins. *Plant Physiol.* 140:311-325. Pubmed:16377746.

- Zhu, J; Mickelson, SM; Kaepller, SM; Lynch, JP, 2006. Detection of quantitative trait loci for seminal root traits in maize (*Zea mays* L.) seedlings grown under differential phosphorus levels. *Theor. Appl. Genet.* 113:1-10. Pubmed:16783587.
- Zimmermann, R; Werr, W, 2007. Transcription of the putative maize orthologue of the *Arabidopsis DORNROSCHEN* gene marks early asymmetry in the proembryo and during leaf initiation in the shoot apical meristem. *Gene Expr. Patterns* 7:158-164. Pubmed:16880009.